

Mouse Dyrk3 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP14712a**Specification**

Mouse Dyrk3 Antibody (N-term) Blocking Peptide - Product InformationPrimary Accession [Q922Y0](#)**Mouse Dyrk3 Antibody (N-term) Blocking Peptide - Additional Information****Gene ID** 226419**Other Names**Dual specificity tyrosine-phosphorylation-regulated kinase 3, Dyrk3
{ECO:0000312|MGI:MGI:1330300}**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Mouse Dyrk3 Antibody (N-term) Blocking Peptide - Protein Information**Name** Dyrk3 {ECO:0000312|MGI:MGI:1330300}**Function**

Dual-specificity protein kinase that promotes disassembly of several types of membraneless organelles during mitosis, such as stress granules, nuclear speckles and pericentriolar material (By similarity). Dual-specificity tyrosine-regulated kinases (DYRKs) autophosphorylate a critical tyrosine residue in their activation loop and phosphorylate their substrate on serine and threonine residues (PubMed:12356771). Acts as a central dissolvase of membraneless organelles during the G2- to-M transition, after the nuclear-envelope breakdown: acts by mediating phosphorylation of multiple serine and threonine residues in unstructured domains of proteins, such as SRRM1 and PCM1 (By similarity). Does not mediate disassembly of all membraneless organelles: disassembly of P-body and nucleolus is not regulated by DYRK3 (By similarity). Dissolution of membraneless organelles at the onset of mitosis is also required to release mitotic regulators, such as ZNF207, from liquid-unmixed organelles where they are sequestered and keep them dissolved during mitosis (By similarity). Regulates mTORC1 by mediating the dissolution of stress granules: during stressful conditions, DYRK3 partitions from the cytosol to the stress granule, together with mTORC1 components, which prevents mTORC1 signaling (By similarity). When stress signals are gone, the kinase activity of DYRK3 is required for the dissolution of stress granule and mTORC1 relocation to the cytosol: acts by mediating the phosphorylation of the mTORC1 inhibitor AKT1S1,

allowing full reactivation of mTORC1 signaling (By similarity). Also acts as a negative regulator of EPO- dependent erythropoiesis: may place an upper limit on red cell production during stress erythropoiesis (By similarity). Inhibits cell death due to cytokine withdrawal in hematopoietic progenitor cells (By similarity). Promotes cell survival upon genotoxic stress through phosphorylation of SIRT1: this in turn inhibits p53/TP53 activity and apoptosis (PubMed:20167603).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:O43781}. Cytoplasm {ECO:0000250|UniProtKB:O43781}. Nucleus speckle {ECO:0000250|UniProtKB:O43781}. Cytoplasmic granule {ECO:0000250|UniProtKB:O43781}. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome {ECO:0000250|UniProtKB:O43781} Note=Associates with membraneless organelles in the cytoplasm and nucleus. Shuttles between cytoplasm and stress granules. Localized predominantly on distinct speckles distributed throughout the cytoplasm of the cell. At low concentration, shows a homogeneous distribution throughout the cytoplasm and does not condense in speckles. During oxidative and osmotic stress, localizes to stress granules {ECO:0000250|UniProtKB:O43781}

Mouse Dyrk3 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

Mouse Dyrk3 Antibody (N-term) Blocking Peptide - Images**Mouse Dyrk3 Antibody (N-term) Blocking Peptide - Background**

Negative regulator of EPO-dependent erythropoiesis, may place an upper limit on red cell production during stress erythropoiesis. Inhibits cell death due to cytokine withdrawal in hematopoietic progenitor cells. May act by regulating CREB/CRE signaling.

Mouse Dyrk3 Antibody (N-term) Blocking Peptide - References

Guo, X., et al. J. Biol. Chem. 285(17):13223-13232(2010)Bogacheva, O., et al. J. Biol. Chem. 283(52):36665-36675(2008)Wojchowski, D.M., et al. Blood Cells Mol. Dis. 36(2):232-238(2006)Zhang, D., et al. Genomics 85(1):117-130(2005)Li, K., et al. J. Biol. Chem. 277(49):47052-47060(2002)