

## **CDH7 Antibody (C-term) Blocking Peptide** Synthetic peptide

Catalog # BP1471b

## Specification

## CDH7 Antibody (C-term) Blocking Peptide - Product Information

Primary Accession

<u>Q9ULB5</u>

## CDH7 Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 1005

Other Names Cadherin-7, CDH7, CDH7L1

Target/Specificity

The synthetic peptide sequence used to generate the antibody <a href=/product/products/AP1471b>AP1471b</a> was selected from the C-term region of human CDH7. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage** Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### **Precautions** This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## CDH7 Antibody (C-term) Blocking Peptide - Protein Information

Name CDH7

Synonyms CDH7L1

Function

Cadherins are calcium-dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types.

**Cellular Location** 

Cell membrane; Single-pass type I membrane protein

# CDH7 Antibody (C-term) Blocking Peptide - Protocols



Provided below are standard protocols that you may find useful for product applications.

#### Blocking Peptides

### CDH7 Antibody (C-term) Blocking Peptide - Images

### CDH7 Antibody (C-term) Blocking Peptide - Background

CDH7 is a type II classical cadherin from the cadherin superfamily. This protein is a calcium dependent cell-cell adhesion glycoprotein comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. Type II (atypical) cadherins are defined based on their lack of a HAV cell adhesion recognition sequence specific to type I cadherins. Cadherins mediate cell-cell binding in a homophilic manner, contributing to the sorting of heterogeneous cell types and the maintenance of orderly structures.

### CDH7 Antibody (C-term) Blocking Peptide - References

Moore,R., Oncogene 23 (40), 6726-6735 (2004)Kools,P., Genomics 68 (3), 283-295 (2000)Shimoyama,Y., Biochem. J. 349 (PT 1), 159-167 (2000)Kremmidiotis,G., Genomics 49 (3), 467-471 (1998)