

PKD2 Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP14827b**Specification**

PKD2 Antibody (C-term) Blocking Peptide - Product InformationPrimary Accession [Q13563](#)**PKD2 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 5311**Other Names**

Polycystin-2, Autosomal dominant polycystic kidney disease type II protein, Polycystic kidney disease 2 protein, Polycystin, R48321, PKD2

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

PKD2 Antibody (C-term) Blocking Peptide - Protein Information**Name** PKD2 ([HGNC:9009](#))**Function**

Component of a heteromeric calcium-permeable ion channel formed by PKD1 and PKD2 that is activated by interaction between PKD1 and a Wnt family member, such as WNT3A and WNT9B (PubMed: [27214281](http://www.uniprot.org/citations/27214281)). Can also form a functional, homotetrameric ion channel (PubMed: [29899465](http://www.uniprot.org/citations/29899465)). Functions as a cation channel involved in fluid-flow mechanosensation by the primary cilium in renal epithelium (PubMed: [18695040](http://www.uniprot.org/citations/18695040)). Functions as outward-rectifying K(+) channel, but is also permeable to Ca(2+), and to a much lesser degree also to Na(+) (PubMed: [11854751](http://www.uniprot.org/citations/11854751), PubMed: [15692563](http://www.uniprot.org/citations/15692563), PubMed: [27071085](http://www.uniprot.org/citations/27071085), PubMed: [27991905](http://www.uniprot.org/citations/27991905)). May contribute to the release of Ca(2+) stores from the endoplasmic reticulum (PubMed: [11854751](http://www.uniprot.org/citations/11854751), PubMed: [20881056](http://www.uniprot.org/citations/20881056)). Together with TRPV4, forms mechano- and thermosensitive channels in cilium (PubMed: [18695040](http://www.uniprot.org/citations/18695040)).

target="_blank">18695040). PKD1 and PKD2 may function through a common signaling pathway that is necessary to maintain the normal, differentiated state of renal tubule cells. Acts as a regulator of cilium length, together with PKD1. The dynamic control of cilium length is essential in the regulation of mechanotransductive signaling. The cilium length response creates a negative feedback loop whereby fluid shear-mediated deflection of the primary cilium, which decreases intracellular cAMP, leads to cilium shortening and thus decreases flow-induced signaling. Also involved in left-right axis specification via its role in sensing nodal flow; forms a complex with PKD1L1 in cilia to facilitate flow detection in left- right patterning. Detection of asymmetric nodal flow gives rise to a Ca(2+) signal that is required for normal, asymmetric expression of genes involved in the specification of body left-right laterality (By similarity).

Cellular Location

Cell projection, cilium membrane; Multi-pass membrane protein. Endoplasmic reticulum membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Basolateral cell membrane. Cytoplasmic vesicle membrane. Golgi apparatus {ECO:0000250|UniProtKB:O35245}. Vesicle Secreted, extracellular exosome Note=PKD2 localization to the plasma and ciliary membranes requires PKD1. PKD1:PKD2 interaction is required to reach the Golgi apparatus form endoplasmic reticulum and then traffic to the cilia (By similarity). Retained in the endoplasmic reticulum by interaction with PACS1 and PACS2 (PubMed:15692563). Detected on kidney tubule basolateral membranes and basal cytoplasmic vesicles (PubMed:10770959) Cell surface and cilium localization requires GANAB (PubMed:27259053) Detected on migrasomes and on extracellular exosomes in urine (PubMed:21406692). {ECO:0000250|UniProtKB:O35245, ECO:0000269|PubMed:15692563, ECO:0000269|PubMed:21406692, ECO:0000269|PubMed:27259053}

Tissue Location

Detected in fetal and adult kidney (PubMed:10770959). Detected at the thick ascending limb of the loop of Henle, at distal tubules, including the distal convoluted tubule and cortical collecting tubules, with weak staining of the collecting duct (PubMed:10770959). Detected on placenta syncytiotrophoblasts (at protein level) (PubMed:26269590). Strongly expressed in ovary, fetal and adult kidney, testis, and small intestine. Not detected in peripheral leukocytes.

PKD2 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

PKD2 Antibody (C-term) Blocking Peptide - Images

PKD2 Antibody (C-term) Blocking Peptide - Background

This gene encodes a member of the polycystin protein family. The encoded protein contains multiple transmembrane domains, and cytoplasmic N- and C-termini. The protein may be an integral membrane protein involved in cell-cell/matrix interactions. The encoded protein may function in renal tubular development, morphology, and function, and may modulate intracellular calcium homeostasis and other signal transduction pathways. This protein interacts with polycystin 1 to produce cation-permeable currents. Mutations in this gene have been associated with autosomal dominant polycystic kidney disease.

PKD2 Antibody (C-term) Blocking Peptide - References

Duning, K., et al. J. Biol. Chem. 285(44):33584-33588(2010) Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010) Stewart, A.P., et al. Biophys. J. 99(3):790-797(2010) Petri, E.T., et al. Proc. Natl. Acad. Sci. U.S.A. 107(20):9176-9181(2010) Steiner, T.S., et al. Cell. Immunol. 264(2):135-142(2010)