

SSRP1 Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP14831b**Specification**

SSRP1 Antibody (C-term) Blocking Peptide - Product InformationPrimary Accession [Q08945](#)**SSRP1 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 6749**Other Names**

FACT complex subunit SSRP1, Chromatin-specific transcription elongation factor 80 kDa subunit, Facilitates chromatin transcription complex 80 kDa subunit, FACT 80 kDa subunit, FACTp80, Facilitates chromatin transcription complex subunit SSRP1, Recombination signal sequence recognition protein 1, Structure-specific recognition protein 1, hSSRP1, T160, SSRP1, FACT80

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

SSRP1 Antibody (C-term) Blocking Peptide - Protein Information**Name** SSRP1**Synonyms** FACT80**Function**

Component of the FACT complex, a general chromatin factor that acts to reorganize nucleosomes. The FACT complex is involved in multiple processes that require DNA as a template such as mRNA elongation, DNA replication and DNA repair. During transcription elongation the FACT complex acts as a histone chaperone that both destabilizes and restores nucleosomal structure. It facilitates the passage of RNA polymerase II and transcription by promoting the dissociation of one histone H2A-H2B dimer from the nucleosome, then subsequently promotes the reestablishment of the nucleosome following the passage of RNA polymerase II. The FACT complex is probably also involved in phosphorylation of 'Ser-392' of p53/TP53 via its association with CK2 (casein kinase II). Binds specifically to double-stranded DNA and at low levels to DNA modified by the antitumor agent cisplatin. May potentiate cisplatin-induced cell death by blocking replication and repair of modified DNA. Also acts as a transcriptional coactivator for p63/TP63.

Cellular Location

Nucleus. Nucleus, nucleolus. Chromosome. Note=Colocalizes with RNA polymerase II on chromatin. Recruited to actively transcribed loci {ECO:0000250|UniProtKB:Q05344}

SSRP1 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

SSRP1 Antibody (C-term) Blocking Peptide - Images

SSRP1 Antibody (C-term) Blocking Peptide - Background

The protein encoded by this gene is a subunit of a heterodimer that, along with SUPT16H, forms chromatin transcriptional elongation factor FACT. FACT interacts specifically with histones H2A/H2B to effect nucleosome disassembly and transcription elongation. FACT and cisplatin-damaged DNA may be crucial to the anticancer mechanism of cisplatin. This encoded protein contains a high mobility group box which most likely constitutes the structure recognition element for cisplatin-modified DNA. This protein also functions as a co-activator of the transcriptional activator p63. An alternatively spliced transcript variant of this gene has been described, but its full-length nature is not known.

SSRP1 Antibody (C-term) Blocking Peptide - References

Zeng, S.X., et al. Mol. Cell. Biol. 30(4):935-947(2010) Hautbergue, G.M., et al. Curr. Biol. 19(22):1918-1924(2009) Hu, J., et al. J. Virol. 83(21):11051-11063(2009) Kumari, A., et al. J. Cell. Biochem. 108(2):508-518(2009) Okada, M., et al. Mol. Biol. Cell 20(18):3986-3995(2009)