10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

NANOG Antibody (S285) Blocking Peptide
Synthetic peptide
Catalog # BP1486e

Specification

NANOG Antibody (S285) Blocking Peptide - Product Information

Primary Accession	Q9H9S0
Other Accession	NP_079141

NANOG Antibody (S285) Blocking Peptide - Additional Information

Gene ID 79923

Other Names
Homeobox protein NANOG, Homeobox transcription factor Nanog, hNanog, NANOG

Target/Specificity
The synthetic peptide sequence used to generate the antibody AP1486e was selected from the S285 region of human NANOG. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format
Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage
Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions
This product is for research use only. Not for use in diagnostic or therapeutic procedures.

NANOG Antibody (S285) Blocking Peptide - Protein Information

Name NANOG

Function
Transcription regulator involved in inner cell mass and embryonic stem (ES) cells proliferation and self-renewal. Imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and trophectoderm lineages. Blocks bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators to the active SMAD transcriptional complexes. Acts as a transcriptional activator or repressor. Binds optimally to the DNA consensus sequence 5'-TAAT[GT][GT]-3' or 5'-[CG][GA][CG]C[GC]ATTAN[GC]- 3'. Binds to the POU5F1/OCT4 promoter (PubMed:25825768). Able to autorepress its expression in differentiating (ES) cells: binds to its own promoter following interaction with ZNF281/ZFP281, leading to recruitment of the NuRD complex and subsequent repression of expression. When overexpressed, promotes cells to enter into S phase and

proliferation.

Cellular Location
Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108, ECO:0000269|PubMed:15983365}

Tissue Location
Expressed in testicular carcinoma and derived germ cell tumors (at protein level). Expressed in fetal gonads, ovary and testis. Also expressed in ovary teratocarcinoma cell line and testicular embryonic carcinoma. Not expressed in many somatic organs and oocytes.

NANOG Antibody (S285) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Blocking Peptides

NANOG Antibody (S285) Blocking Peptide - Images

NANOG Antibody (S285) Blocking Peptide - Background

NANOG is a Ttranscription regulator involved in inner cell mass and embryonic stem (ES) cels proliferation and self-renewal. It imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and trophectoderm lineages. This protein blocks bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators to the active SMAD transcriptional complexes. NANOG acts as a transcriptional activator or repressor. It binds optimally to the DNA consensus sequence 5'-[CG][GA][CG]C[GC]ATTAN[GC]-3'. When overexpressed, this protein promotes cells to enter into S phase and proliferation.

NANOG Antibody (S285) Blocking Peptide - References

Kochupurakkal,B.S., Biochem. Biophys. Res. Commun. 365 (4), 846-850 (2008)Freberg,C.T., Mol. Biol. Cell 18 (5), 1543-1553 (2007)