

RBM22 Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP14906c**Specification**

RBM22 Antibody (Center) Blocking Peptide - Product InformationPrimary Accession [Q9NW64](#)**RBM22 Antibody (Center) Blocking Peptide - Additional Information****Gene ID** 55696**Other Names**

Pre-mRNA-splicing factor RBM22, RNA-binding motif protein 22, Zinc finger CCCH domain-containing protein 16, RBM22, ZC3H16

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

RBM22 Antibody (Center) Blocking Peptide - Protein Information**Name** RBM22**Synonyms** ZC3H16**Function**

Required for pre-mRNA splicing as component of the activated spliceosome (PubMed: 28502770, PubMed: 28076346, PubMed: 29361316, PubMed: 29360106, PubMed: 29301961, PubMed: 30705154). Involved in the first step of pre-mRNA splicing. Binds directly to the internal stem-loop (ISL) domain of the U6 snRNA and to the pre-mRNA intron near the 5' splice site during the activation and catalytic phases of the spliceosome cycle. Involved in both translocations of the nuclear SLU7 to the cytoplasm and the cytosolic calcium-binding protein PDCD6 to the nucleus upon cellular stress responses.

Cellular Location

Nucleus. Cytoplasm Note=Nearly exclusively nuclear. Translocated from the nucleus to the

cytoplasm after heat shock cell treatment. May be shuttling between the nucleus and the cytosol.

RBM22 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

RBM22 Antibody (Center) Blocking Peptide - Images

RBM22 Antibody (Center) Blocking Peptide - Background

This gene encodes an RNA binding protein. The encoded protein may play a role in cell division and may be involved in pre-mRNA splicing. Related pseudogenes exist on chromosomes 6, 7, 9, 13, 16, 18, and X.

RBM22 Antibody (Center) Blocking Peptide - References

Krebs, J. Biochim. Biophys. Acta 1793(6):979-984(2009) Satoh, J., et al. Neuropathol. Appl. Neurobiol. 35(1):16-35(2009) He, F., et al. Genet. Mol. Res. 8(4):1466-1473(2009) Montaville, P., et al. Biochim. Biophys. Acta 1763(11):1335-1343(2006) Kittler, R., et al. Nature 432(7020):1036-1040(2004)