

**Mouse Pak1 Antibody (C-term) Blocking Peptide**  
Synthetic peptide  
Catalog # BP14922b**Specification**

---

**Mouse Pak1 Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [O88643](#)**Mouse Pak1 Antibody (C-term) Blocking Peptide - Additional Information****Other Names**

Serine/threonine-protein kinase PAK 1, Alpha-PAK, CDC42/RAC effector kinase PAK-A, p21-activated kinase 1, PAK-1, p65-PAK, Pak1, Paka

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**Mouse Pak1 Antibody (C-term) Blocking Peptide - Protein Information**

Name Pak1 {ECO:0000312|MGI:MGI:1339975}

**Synonyms** Paka**Function**

Protein kinase involved in intracellular signaling pathways downstream of integrins and receptor-type kinases that plays an important role in cytoskeleton dynamics, in cell adhesion, migration, proliferation, apoptosis, mitosis, and in vesicle-mediated transport processes (PubMed: [10611223](http://www.uniprot.org/citations/10611223)), PubMed: [12176334](http://www.uniprot.org/citations/12176334), PubMed: [12165471](http://www.uniprot.org/citations/12165471), PubMed: [15800193](http://www.uniprot.org/citations/15800193), PubMed: [22669945](http://www.uniprot.org/citations/22669945), PubMed: [23633677](http://www.uniprot.org/citations/23633677)). Can directly phosphorylate BAD and protects cells against apoptosis (PubMed: [10611223](http://www.uniprot.org/citations/10611223)). Activated by interaction with CDC42 and RAC1 (By similarity). Functions as a GTPase effector that links the Rho-related GTPases CDC42 and RAC1 to the JNK MAP kinase pathway (By similarity). Phosphorylates and activates MAP2K1, and thereby mediates activation of downstream MAP kinases (By similarity). Involved in the reorganization of the actin cytoskeleton, actin stress fibers and of focal adhesion complexes (By similarity). Phosphorylates the tubulin chaperone TBCB and thereby plays a role in the regulation of microtubule biogenesis and organization of the tubulin

cytoskeleton (PubMed: <a href="http://www.uniprot.org/citations/12176334" target="\_blank">12176334</a>). Plays a role in the regulation of insulin secretion in response to elevated glucose levels (PubMed: <a href="http://www.uniprot.org/citations/22669945" target="\_blank">22669945</a>). Part of a ternary complex that contains PAK1, DVL1 and MUSK that is important for MUSK-dependent regulation of AChR clustering during the formation of the neuromuscular junction (NMJ) (PubMed: <a href="http://www.uniprot.org/citations/12165471" target="\_blank">12165471</a>). Activity is inhibited in cells undergoing apoptosis, potentially due to binding of CDC2L1 and CDC2L2 (By similarity). Phosphorylates MYL9/MLC2 (By similarity). Phosphorylates RAF1 at 'Ser-338' and 'Ser-339' resulting in: activation of RAF1, stimulation of RAF1 translocation to mitochondria, phosphorylation of BAD by RAF1, and RAF1 binding to BCL2 (By similarity). Phosphorylates SNAIL1 at 'Ser-246' promoting its transcriptional repressor activity by increasing its accumulation in the nucleus (By similarity). In podocytes, promotes NR3C2 nuclear localization (By similarity). Required for atypical chemokine receptor ACKR2-induced phosphorylation of LIMK1 and cofilin (CFL1) and for the up-regulation of ACKR2 from endosomal compartment to cell membrane, increasing its efficiency in chemokine uptake and degradation (PubMed: <a href="http://www.uniprot.org/citations/23633677" target="\_blank">23633677</a>). In synapses, seems to mediate the regulation of F-actin cluster formation performed by SHANK3, maybe through CFL1 phosphorylation and inactivation (By similarity). Plays a role in RUFY3-mediated facilitating gastric cancer cells migration and invasion (By similarity). In response to DNA damage, phosphorylates MORC2 which activates its ATPase activity and facilitates chromatin remodeling (By similarity). In neurons, plays a crucial role in regulating GABA(A) receptor synaptic stability and hence GABAergic inhibitory synaptic transmission through its role in F-actin stabilization (By similarity). In hippocampal neurons, necessary for the formation of dendritic spines and excitatory synapses; this function is dependent on kinase activity and may be exerted by the regulation of actomyosin contractility through the phosphorylation of myosin II regulatory light chain (MLC) (PubMed: <a href="http://www.uniprot.org/citations/15800193" target="\_blank">15800193</a>). Along with GIT1, positively regulates microtubule nucleation during interphase (By similarity). Phosphorylates FXR1, promoting its localization to stress granules and activity (By similarity).

#### Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q13153}. Cell junction, focal adhesion {ECO:0000250|UniProtKB:Q13153}. Cell projection, lamellipodium {ECO:0000250|UniProtKB:Q13153}. Cell membrane {ECO:0000250|UniProtKB:Q13153}. Cell projection, ruffle membrane {ECO:0000250|UniProtKB:Q13153}. Cell projection, invadopodium {ECO:0000250|UniProtKB:Q13153}. Nucleus, nucleoplasm {ECO:0000250|UniProtKB:Q13153}. Chromosome {ECO:0000250|UniProtKB:Q13153}. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome {ECO:0000250|UniProtKB:Q13153} Note=Recruited to the cell membrane by interaction with CDC42 and RAC1 (By similarity). Recruited to focal adhesions upon activation Colocalized with CIB1 within membrane ruffles during cell spreading upon readhesion to fibronectin. Colocalizes with RUFY3, F-actin and other core migration components in invadopodia at the cell periphery (By similarity). Upon DNA damage, translocates to the nucleoplasm when phosphorylated at Thr-212 where is co-recruited with MORC2 on damaged chromatin (By similarity). Localization to the centrosome does not depend upon the presence of gamma-tubulin (By similarity). Localization of the active, but not inactive, protein to the adhesions and edge of lamellipodia is mediated by interaction with GIT1 (By similarity) {ECO:0000250|UniProtKB:P35465, ECO:0000250|UniProtKB:Q13153}

#### Mouse Pak1 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

#### Mouse Pak1 Antibody (C-term) Blocking Peptide - Images

#### Mouse Pak1 Antibody (C-term) Blocking Peptide - Background

The activated kinase acts on a variety of targets. Likely to be the GTPase effector that links the Rho-related GTPases to the JNK MAP kinase pathway. Activated by CDC42 and RAC1. Involved in dissolution of stress fibers and reorganization of focal complexes. Involved in regulation of microtubule biogenesis through phosphorylation of TBCB. Activity is inhibited in cells undergoing apoptosis, potentially due to binding of CDC2L1 and CDC2L2 (By similarity).