

SAP18 Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP14996c**Specification**

SAP18 Antibody (Center) Blocking Peptide - Product InformationPrimary Accession [O00422](#)**SAP18 Antibody (Center) Blocking Peptide - Additional Information****Gene ID** 10284**Other Names**

Histone deacetylase complex subunit SAP18, 18 kDa Sin3-associated polypeptide, 2HOR0202, Cell growth-inhibiting gene 38 protein, Sin3-associated polypeptide p18, SAP18

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

SAP18 Antibody (Center) Blocking Peptide - Protein Information**Name** SAP18**Function**

Component of the SIN3-repressing complex. Enhances the ability of SIN3-HDAC1-mediated transcriptional repression. When tethered to the promoter, it can direct the formation of a repressive complex to core histone proteins. Auxiliary component of the splicing- dependent multiprotein exon junction complex (EJC) deposited at splice junction on mRNAs. The EJC is a dynamic structure consisting of core proteins and several peripheral nuclear and cytoplasmic associated factors that join the complex only transiently either during EJC assembly or during subsequent mRNA metabolism. Component of the ASAP and PSAP complexes which bind RNA in a sequence-independent manner and are proposed to be recruited to the EJC prior to or during the splicing process and to regulate specific excision of introns in specific transcription subsets. The ASAP complex can inhibit mRNA processing during in vitro splicing reactions. The ASAP complex promotes apoptosis and is disassembled after induction of apoptosis. Involved in the splicing modulation of BCL2L1/Bcl-X (and probably other apoptotic genes); specifically inhibits the formation of proapoptotic isoforms such as Bcl-X(S); the activity is different from the established EJC assembly and function.

Cellular Location

Nucleus. Cytoplasm. Nucleus speckle. Note=Shuttles between the nucleus and the cytoplasm

(PubMed:16314458). Colocalizes with ACIN1 and SRSF2 in nuclear speckles (PubMed:20966198).

Tissue Location

Ubiquitous.

SAP18 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

SAP18 Antibody (Center) Blocking Peptide - Images**SAP18 Antibody (Center) Blocking Peptide - Background**

Histone acetylation plays a key role in the regulation of eukaryotic gene expression. Histone acetylation and deacetylation are catalyzed by multisubunit complexes. The protein encoded by this gene is a component of the histone deacetylase complex, which includes SIN3, SAP30, HDAC1, HDAC2, RbAp46, RbAp48, and other polypeptides. This protein directly interacts with SIN3 and enhances SIN3-mediated transcriptional repression when tethered to the promoter. A pseudogene has been identified on chromosome 2.

SAP18 Antibody (Center) Blocking Peptide - References

Sorin, M., et al. PLoS Pathog. 5 (6), E1000463 (2009) ; McCallum, S.A., et al. Biochemistry 45(39):11974-11982(2006) ; Joselin, A.P., et al. J. Biol. Chem. 281(18):12475-12484(2006) ; Schwerk, C., et al. Mol. Cell. Biol. 23(8):2981-2990(2003) ; Koipally, J., et al. EMBO J. 18(11):3090-3100(1999)