

### Connexin 36 Antibody (C-term) Blocking peptide

Synthetic peptide Catalog # BP1549b

### **Specification**

### Connexin 36 Antibody (C-term) Blocking peptide - Product Information

**Primary Accession** 

09UKL4

### Connexin 36 Antibody (C-term) Blocking peptide - Additional Information

**Gene ID 57369** 

#### **Other Names**

Gap junction delta-2 protein, Connexin-36, Cx36, Gap junction alpha-9 protein, GID2, GIA9

#### Target/Specificity

The synthetic peptide sequence used to generate the antibody <a href=/product/products/AP1549b>AP1549b</a> was selected from the C-term region of human GJA9. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### Connexin 36 Antibody (C-term) Blocking peptide - Protein Information

Name GJD2

Synonyms GJA9

#### **Function**

One gap junction consists of a cluster of closely packed pairs of transmembrane channels, the connexons, through which materials of low MW diffuse from one cell to a neighboring cell.

#### **Cellular Location**

Cell membrane; Multi-pass membrane protein. Cell junction, gap junction

### **Tissue Location**

Highly expressed in neurons.



# Connexin 36 Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

### • Blocking Peptides

Connexin 36 Antibody (C-term) Blocking peptide - Images

### Connexin 36 Antibody (C-term) Blocking peptide - Background

GJA9, also called connexin-36 (CX36), is a member of the connexin gene family that is expressed predominantly in mammalian neurons. Connexins associate in groups of 6 and are organized radially around a central pore to form connexons. Each gap junction intercellular channel is formed by the conjunction of 2 connexons.

## Connexin 36 Antibody (C-term) Blocking peptide - References

de Brouwer, A.P., et al., Hum. Genet. 112(2):156-163 (2003).Belluardo, N., et al., J. Neurosci. Res. 57(5):740-752 (1999).