

### **NUDC Antibody (Center) Blocking Peptide**

Synthetic peptide Catalog # BP16056c

## **Specification**

## **NUDC Antibody (Center) Blocking Peptide - Product Information**

**Primary Accession** 

**09Y266** 

# **NUDC Antibody (Center) Blocking Peptide - Additional Information**

**Gene ID** 10726

#### **Other Names**

Nuclear migration protein nudC, Nuclear distribution protein C homolog, NUDC

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

# **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## **NUDC Antibody (Center) Blocking Peptide - Protein Information**

## Name NUDC

#### **Function**

Plays a role in neurogenesis and neuronal migration (By similarity). Necessary for correct formation of mitotic spindles and chromosome separation during mitosis (PubMed:<a href="http://www.uniprot.org/citations/12852857" target="\_blank">12852857</a>, PubMed:<a href="http://www.uniprot.org/citations/12679384" target="\_blank">12679384</a>, PubMed:<a href="http://www.uniprot.org/citations/25789526" target="\_blank">25789526</a>). Necessary for cytokinesis and cell proliferation (PubMed:<a

 $href="http://www.uniprot.org/citations/12852857" \ target="\_blank">12852857</a>, PubMed:<a href="http://www.uniprot.org/citations/12679384" \ target="\_blank">12679384</a>).$ 

### **Cellular Location**

Cytoplasm, cytoskeleton. Nucleus. Cytoplasm, cytoskeleton, spindle. Midbody Note=In a filamentous pattern adjacent to the nucleus of migrating cerebellar granule cells. Colocalizes with tubulin and dynein and with the microtubule organizing center. Distributed throughout the cytoplasm of non-migrating cells. A small proportion is nuclear, in a punctate pattern. Localizes to the mitotic spindle in a EML4-dependent manner (PubMed:25789526).

#### **Tissue Location**

Ubiquitous. Highly expressed in fetal liver, kidney, lung and brain. Highly expressed in adult



Tel: 858.875.1900 Fax: 858.875.1999

pancreas, kidney, skeletal muscle, liver, lung, placenta, prostate, brain and heart

# **NUDC Antibody (Center) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

### • Blocking Peptides

**NUDC Antibody (Center) Blocking Peptide - Images** 

# **NUDC Antibody (Center) Blocking Peptide - Background**

NudC was first identified as a regulator of nuclearmovement in the asexual reproductive cycle of the filamentousfungus Aspergillus nidulans. Human NUDC is a nuclear movement that associates with dynein (see DYNC1H1; MIM 600112)(Aumais et al., 2003 [PubMed 12679384]).

### **NUDC Antibody (Center) Blocking Peptide - References**

Zhu, X.J., et al. J. Biol. Chem. 285(39):29903-29910(2010)Chen, W.M., et al. J. Biol. Chem. 285(34):26697-26709(2010)Pang, S.F., et al. Exp. Cell Res. 315(20):3563-3573(2009)Tang, Y.S., et al. Leukemia 22(5):1018-1025(2008)Zhang, Y.P., et al. Exp. Cell Res. 313(15):3210-3221(2007)