

ASGR1 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP16133a**Specification**

ASGR1 Antibody (N-term) Blocking Peptide - Product InformationPrimary Accession [P07306](#)**ASGR1 Antibody (N-term) Blocking Peptide - Additional Information****Gene ID** 432**Other Names**

Asialoglycoprotein receptor 1, ASGP-R 1, ASGPR 1, C-type lectin domain family 4 member H1, Hepatic lectin H1, HL-1, ASGR1, CLEC4H1

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

ASGR1 Antibody (N-term) Blocking Peptide - Protein Information**Name** ASGR1**Synonyms** CLEC4H1**Function**

Mediates the endocytosis of plasma glycoproteins to which the terminal sialic acid residue on their complex carbohydrate moieties has been removed. The receptor recognizes terminal galactose and N- acetylgalactosamine units. After ligand binding to the receptor, the resulting complex is internalized and transported to a sorting organelle, where receptor and ligand are disassociated. The receptor then returns to the cell membrane surface.

Cellular Location

[Isoform H1a]: Membrane; Single-pass type II membrane protein

Tissue Location

Expressed exclusively in hepatic parenchymal cells.

ASGR1 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

ASGR1 Antibody (N-term) Blocking Peptide - Images

ASGR1 Antibody (N-term) Blocking Peptide - Background

Partially deglycosylated plasma glycoproteins and immunoglobulin IgA2 allotypes are efficiently and specifically removed from circulation by a receptor-mediated process. The asialoglycoprotein receptor binds to desialylated (galactosyl-terminal) glycoproteins. It transports these glycoproteins via a series of membrane vesicles and tubules to an acidic-sorting organelle where the receptor and ligand dissociate. Then the receptor is recycled back to the cell surface and the ligand is transported to the lysosomes for degradation. Alternatively spliced transcript variants encoding distinct isoforms have been identified.

ASGR1 Antibody (N-term) Blocking Peptide - References

Yang, J., et al. Arch. Virol. 155(6):881-888(2010) Liu, J., et al. PLoS ONE 5 (9), E12934 (2010)
:Sorensen, A.L., et al. Blood 114(8):1645-1654(2009) Oh, J.H., et al. Mamm. Genome
16(12):942-954(2005) Yik, J.H., et al. J. Biol. Chem. 277(43):40844-40852(2002)