

### **CEP72 Antibody (Center) Blocking Peptide** Synthetic peptide

Catalog # BP16441c

## Specification

# **CEP72** Antibody (Center) Blocking Peptide - Product Information

Primary Accession

<u>Q9P209</u>

### **CEP72** Antibody (Center) Blocking Peptide - Additional Information

Gene ID 55722

Other Names Centrosomal protein of 72 kDa, Cep72, CEP72, KIAA1519

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage** Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions** This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### **CEP72** Antibody (Center) Blocking Peptide - Protein Information

Name CEP72

Synonyms KIAA1519

#### Function

Involved in the recruitment of key centrosomal proteins to the centrosome. Provides centrosomal microtubule-nucleation activity on the gamma-tubulin ring complexes (gamma-TuRCs) and has critical roles in forming a focused bipolar spindle, which is needed for proper tension generation between sister chromatids. Required for localization of KIZ, AKAP9 and gamma-tubulin ring complexes (gamma-TuRCs) (PubMed:<a href="http://www.uniprot.org/citations/19536135" target="\_blank">19536135</a>). Involved in centriole duplication. Required for CDK5RAP22, CEP152, WDR62 and CEP63 centrosomal localization and promotes the centrosomal localization of CDK2 (PubMed:<a href="http://www.uniprot.org/citations/26297806" target=" blank">26297806</a>).

#### **Cellular Location**

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriolar satellite. Note=Localizes to the centrosome and centrosome-surrounding particles throughout the cell cycle. These particles disappear after microtubules are depolymerized using nocodazole, suggesting that CEP72-associating particles localize in a microtubule- dependent manner



### **CEP72 Antibody (Center) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

<u>Blocking Peptides</u>

## CEP72 Antibody (Center) Blocking Peptide - Images

#### **CEP72** Antibody (Center) Blocking Peptide - Background

CEP72 is a member of theleucine-rich-repeat (LRR) superfamily of proteins. The protein islocalized to the centrosome, a non-membraneous organelle thatfunctions as the major microtubule-organizing center in animalcells.

#### **CEP72 Antibody (Center) Blocking Peptide - References**

McGovern, D.P., et al. Nat. Genet. 42(4):332-337(2010)Briggs, F.B., et al. Genes Immun. 11(3):199-208(2010)Oshimori, N., et al. EMBO J. 28(14):2066-2076(2009)Xin, X., et al. Genome Res. 19(7):1262-1269(2009)Olsen, J.V., et al. Cell 127(3):635-648(2006)