

PRODH2 Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP16627c**Specification**

PRODH2 Antibody (Center) Blocking Peptide - Product InformationPrimary Accession [Q9UF12](#)**PRODH2 Antibody (Center) Blocking Peptide - Additional Information****Gene ID** 58510**Other Names**

Probable proline dehydrogenase 2, Kidney and liver proline oxidase 1, HsPOX1, Probable proline oxidase 2, PRODH2

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

PRODH2 Antibody (Center) Blocking Peptide - Protein Information**Name** PRODH2 ([HGNC:17325](#))**Function**

Dehydrogenase that converts trans-4-L-hydroxyproline to delta-1-pyrroline-3-hydroxy-5-carboxylate (Hyp) using ubiquinone-10 as the terminal electron acceptor. Can also use proline as a substrate but with a very much lower efficiency. Does not react with other diastereomers of Hyp: trans-4-D-hydroxyproline and cis-4-L- hydroxyproline. Ubiquinone analogs such as menadione, duroquinone and ubiquinone-1 react more efficiently than oxygen as the terminal electron acceptor during catalysis.

PRODH2 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

PRODH2 Antibody (Center) Blocking Peptide - Images**PRODH2 Antibody (Center) Blocking Peptide - Background**

The protein encoded by this gene is similar to prolinedehydrogenase (oxidase) 1, a mitochondrial enzyme which catalyzes the first step in proline catabolism. The function of this protein has not been determined.

PRODH2 Antibody (Center) Blocking Peptide - References

Cooper, S.K., et al. J. Biol. Chem. 283(16):10485-10492(2008)