

NF kappa B Antibody (C-term S536) Blocking Peptide
Synthetic peptide
Catalog # BP16870b**Specification**

NF kappa B Antibody (C-term S536) Blocking Peptide - Product InformationPrimary Accession [Q04206](#)**NF kappa B Antibody (C-term S536) Blocking Peptide - Additional Information****Gene ID** 5970**Other Names**

Transcription factor p65, Nuclear factor NF-kappa-B p65 subunit, Nuclear factor of kappa light polypeptide gene enhancer in B-cells 3, RELA, NFKB3

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

NF kappa B Antibody (C-term S536) Blocking Peptide - Protein Information**Name** RELA**Synonyms** NFKB3**Function**

NF-kappa-B is a pleiotropic transcription factor present in almost all cell types and is the endpoint of a series of signal transduction events that are initiated by a vast array of stimuli related to many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain- containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52. The heterodimeric RELA-NFKB1 complex appears to be most abundant one. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct preferences for different kappa-B sites that they can bind with distinguishable affinity and specificity. Different dimer combinations act as transcriptional activators or repressors, respectively. The NF-kappa-B heterodimeric RELA-NFKB1 and RELA-REL complexes, for instance, function as transcriptional activators. NF-kappa-B is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NF-kappa-B complexes are held in the cytoplasm in an inactive state complexed with members of the NF-kappa-B inhibitor (I- kappa-B) family. In a conventional activation pathway, I-kappa-B is phosphorylated by I-kappa-B kinases (IKKs) in

response to different activators, subsequently degraded thus liberating the active NF-kappa-B complex which translocates to the nucleus. The inhibitory effect of I- kappa-B on NF-kappa-B through retention in the cytoplasm is exerted primarily through the interaction with RELA. RELA shows a weak DNA- binding site which could contribute directly to DNA binding in the NF- kappa-B complex. Beside its activity as a direct transcriptional activator, it is also able to modulate promoters accessibility to transcription factors and thereby indirectly regulate gene expression. Associates with chromatin at the NF-kappa-B promoter region via association with DDX1. Essential for cytokine gene expression in T- cells (PubMed:15790681). The NF-kappa-B homodimeric RELA-RELA complex appears to be involved in invasin-mediated activation of IL-8 expression. Key transcription factor regulating the IFN response during SARS-CoV-2 infection (PubMed:33440148).

Cellular Location

Nucleus. Cytoplasm. Note=Nuclear, but also found in the cytoplasm in an inactive form complexed to an inhibitor (I-kappa-B) (PubMed:1493333). Colocalized with DDX1 in the nucleus upon TNF-alpha induction (PubMed:19058135). Colocalizes with GFI1 in the nucleus after LPS stimulation (PubMed:20547752). Translocation to the nucleus is impaired in L.monocytogenes infection (PubMed:20855622)

NF kappa B Antibody (C-term S536) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

NF kappa B Antibody (C-term S536) Blocking Peptide - Images

NF kappa B Antibody (C-term S536) Blocking Peptide - Background

NFKB1 (MIM 164011) or NFKB2 (MIM 164012) is bound to REL(MIM 164910), RELA, or RELB (MIM 604758) to form the NFKB complex.The p50 (NFKB1)/p65 (RELA) heterodimer is the most abundant form ofNFKB. The NFKB complex is inhibited by I-kappa-B proteins (NFKBIA,MIM 164008 or NFKBIB, MIM 604495), which inactivate NFKB bytrapping it in the cytoplasm. Phosphorylation of serine residues onthe I-kappa-B proteins by kinases (IKBKA, MIM 600664, or IKBKB, MIM603258) marks them for destruction via the ubiquitination pathway,thereby allowing activation of the NFKB complex. Activated NFKBcomplex translocates into the nucleus and binds DNA atkappa-B-binding motifs such as 5-prime GGGRNNYYCC 3-prime or5-prime HGGARNYYCC 3-prime (where H is A, C, or T; R is an A or Gpurine; and Y is a C or T pyrimidine).

NF kappa B Antibody (C-term S536) Blocking Peptide - References

Pan, W.W., et al. J. Biol. Chem. 285(45):34348-34354(2010)Tago, K., et al. J. Biol. Chem. 285(40):30622-30633(2010)Park, J.S., et al. Oncol. Rep. 24(3):709-714(2010)Yu, Z.H., et al. Xi Bao Yu Fen Zi Mian Yi Xue Za Zhi 26(7):650-652(2010)Rohwer, N., et al. PLoS ONE 5 (8), E12038 (2010)
: