

## **DUSP12 Antibody (Center) Blocking Peptide**

Synthetic peptide Catalog # BP16903c

# **Specification**

## **DUSP12 Antibody (Center) Blocking Peptide - Product Information**

**Primary Accession** 

**09UNI6** 

# **DUSP12 Antibody (Center) Blocking Peptide - Additional Information**

### **Gene ID 11266**

#### **Other Names**

Dual specificity protein phosphatase 12, Dual specificity tyrosine phosphatase YVH1, DUSP12

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### **DUSP12 Antibody (Center) Blocking Peptide - Protein Information**

## Name DUSP12

### **Function**

Dual specificity phosphatase; can dephosphorylate both phosphotyrosine and phosphoserine or phosphothreonine residues. Can dephosphorylate glucokinase (in vitro) (By similarity). Has phosphatase activity with the synthetic substrate 6,8-difluoro-4-methylumbelliferyl phosphate and other in vitro substrates (PubMed:<a href="http://www.uniprot.org/citations/10446167" target="\_blank">10446167</a>, PubMed:<a href="http://www.uniprot.org/citations/24531476" target="\_blank">24531476</a>).

### **Cellular Location**

Nucleus. Cytoplasm, cytosol. Note=Primarily nuclear. Detected in a mesh-like pattern in the cytosol.

#### **Tissue Location**

Ubiquitous, highest expression in spleen, testis, ovary, and peripheral blood leukocytes and lower expression in liver and lung

## **DUSP12 Antibody (Center) Blocking Peptide - Protocols**



Provided below are standard protocols that you may find useful for product applications.

### Blocking Peptides

## **DUSP12 Antibody (Center) Blocking Peptide - Images**

# **DUSP12 Antibody (Center) Blocking Peptide - Background**

The protein encoded by this gene is a member of the dualspecificity protein phosphatase subfamily. These phosphatasesinactivate their target kinases by dephosphorylating both thephosphoserine/threonine and phosphotyrosine residues. Theynegatively regulate members of the mitogen-activated protein (MAP)kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which is associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissuedistribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This gene product is the human ortholog of the Saccharomyces cerevisiae YVH1 protein tyrosine phosphatase. It is localized predominantly in the nucleus, and is novel in that it contains, and is regulated by a zinc finger domain.

## **DUSP12 Antibody (Center) Blocking Peptide - References**

Biernacki, M.A., et al. Cancer Res. 70(3):906-915(2010)Bonham, C.A., et al. J. Biol. Chem. 284(34):22853-22864(2009)Sharda, P.R., et al. Biochem. J. 418(2):391-401(2009)Gao, F., et al. Zhonghua Yi Xue Za Zhi 88(32):2250-2253(2008)Hasstedt, S.J., et al. Ann. Hum. Genet. 72 (PT 2), 163-169 (2008):