

RNF10 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP17072a**Specification**

RNF10 Antibody (N-term) Blocking Peptide - Product InformationPrimary Accession [Q8N5U6](#)**RNF10 Antibody (N-term) Blocking Peptide - Additional Information**

Gene ID 9921

Other Names

RING finger protein 10, RNF10, KIAA0262, RIE2

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

RNF10 Antibody (N-term) Blocking Peptide - Protein Information**Name** RNF10 {ECO:0000303|PubMed:10697961, ECO:0000312|HGNC:HGNC:10055}**Function**

E3 ubiquitin-protein ligase that catalyzes monoubiquitination of 40S ribosomal proteins RPS2/us5 and RPS3/us3 in response to ribosome stalling (PubMed:34348161, PubMed:34469731). Part of a ribosome quality control that takes place when ribosomes have stalled during translation initiation (iRQC): RNF10 acts by mediating monoubiquitination of RPS2/us5 and RPS3/us3, promoting their degradation by the proteasome (PubMed:34348161, PubMed:34469731). Also promotes ubiquitination of 40S ribosomal proteins in response to ribosome stalling during translation elongation (PubMed:34348161). The action of RNF10 in iRQC is counteracted by USP10 (PubMed:34469731). May also act as a transcriptional factor involved in the regulation of MAG (Myelin-associated glycoprotein) expression (By similarity). Acts as a regulator of Schwann cell differentiation and myelination (By similarity).

Cellular Location

Cytoplasm. Nucleus {ECO:0000250|UniProtKB:Q5XI59}

RNF10 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

RNF10 Antibody (N-term) Blocking Peptide - Images

RNF10 Antibody (N-term) Blocking Peptide - Background

The protein encoded by this gene contains a ring finger motif, which is known to be involved in protein-protein interactions. The specific function of this protein has not yet been determined. EST data suggests the existence of multiple alternatively spliced transcript variants, however, their full length nature is not known.

RNF10 Antibody (N-term) Blocking Peptide - References

Hoshikawa, S., et al. PLoS ONE 3 (10), E3464 (2008) :Stelzl, U., et al. Cell 122(6):957-968(2005) Lin, J., et al. Mol. Cell. Biochem. 275 (1-2), 75-84 (2005) :Seki, N., et al. J. Hum. Genet. 45(1):38-42(2000)