

RIC8A Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP17108c**Specification**

RIC8A Antibody (Center) Blocking Peptide - Product InformationPrimary Accession [Q9NPQ8](#)**RIC8A Antibody (Center) Blocking Peptide - Additional Information****Gene ID** 60626**Other Names**

Synembryn-A, Protein Ric-8A, RIC8A

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

RIC8A Antibody (Center) Blocking Peptide - Protein Information**Name** RIC8A**Function**

Guanine nucleotide exchange factor (GEF), which can activate some, but not all, G-alpha proteins. Able to activate GNAI1, GNAO1 and GNAQ, but not GNAS by exchanging bound GDP for free GTP. Involved in regulation of microtubule pulling forces during mitotic movement of chromosomes by stimulating G(i)-alpha protein, possibly leading to release G(i)-alpha-GTP and NuMA proteins from the NuMA-GPSM2-G(i)-alpha-GDP complex (By similarity). Also acts as an activator for G(q)-alpha (GNAQ) protein by enhancing the G(q)-coupled receptor-mediated ERK activation.

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q80ZG1}. Cell membrane {ECO:0000250|UniProtKB:Q80ZG1}. Note=Colocalizes with RGS14 in CA2 hippocampal neurons. Colocalizes with GNAI1 and RGS14 at the plasma membrane (By similarity).

RIC8A Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

RIC8A Antibody (Center) Blocking Peptide - Images

RIC8A Antibody (Center) Blocking Peptide - Background

Guanine nucleotide exchange factor (GEF), which can activate some, but not all, G-alpha proteins. Able to activate GNAI1, GNAO1 and GNAQ, but not GNAS by exchanging bound GDP for free GTP. Involved in regulation of microtubule pulling forces during mitotic movement of chromosomes by stimulating G(i)-alpha protein, possibly leading to release G(i)-alpha-GTP and NuMA proteins from the NuMA-GPSM2-G(i)-alpha-GDP complex (By similarity). Also acts as an activator for G(q)-alpha (GNAQ) protein by enhancing the G(q)-coupled receptor-mediated ERK activation.

RIC8A Antibody (Center) Blocking Peptide - References

Woodard, G.E., et al. Mol. Cell. Biol. 30(14):3519-3530(2010)Matsuoka, S., et al. Science 316(5828):1160-1166(2007)Lim, J., et al. Cell 125(4):801-814(2006)Nishimura, A., et al. Genes Cells 11(5):487-498(2006)Tall, G.G., et al. J. Biol. Chem. 278(10):8356-8362(2003)