

# **RBM23 Antibody (Center) Blocking Peptide**

Synthetic peptide Catalog # BP17329c

## **Specification**

# **RBM23 Antibody (Center) Blocking Peptide - Product Information**

**Primary Accession** 

**086U06** 

# RBM23 Antibody (Center) Blocking Peptide - Additional Information

**Gene ID 55147** 

### **Other Names**

Probable RNA-binding protein 23, RNA-binding motif protein 23, RNA-binding region-containing protein 4, Splicing factor SF2, RBM23, RNPC4

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

## **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## RBM23 Antibody (Center) Blocking Peptide - Protein Information

Name RBM23 {ECO:0000303|PubMed:31693891, ECO:0000312|HGNC:HGNC:20155}

### **Function**

RNA-binding protein that acts both as a transcription coactivator and pre-mRNA splicing factor (PubMed:<a href="http://www.uniprot.org/citations/15694343" target="\_blank">15694343</a>). Regulates steroid hormone receptor-mediated transcription, independently of the pre-mRNA splicing factor activity (PubMed:<a href="http://www.uniprot.org/citations/15694343" target="\_blank">15694343</a>).

### **Cellular Location**

Nucleus.

### **Tissue Location**

Highly expressed in placenta, liver, skeletal muscle, heart and kidney (PubMed:15694343). Expressed at lower levels in the colon, thymus, spleen, small intestine and lung (PubMed:15694343).

# RBM23 Antibody (Center) Blocking Peptide - Protocols



Provided below are standard protocols that you may find useful for product applications.

## • Blocking Peptides

# **RBM23 Antibody (Center) Blocking Peptide - Images**

# RBM23 Antibody (Center) Blocking Peptide - Background

This gene encodes a member of the U2AF-like family of RNAbinding proteins. This protein interacts with some steroid nuclearreceptors, localizes to the promoter of a steroid- responsive gene, and increases transcription of steroid-responsive transcriptional reporters in a hormone-dependent manner. It is also implicated in the steroid receptor-dependent regulation of alternative splicing. Multiple transcript variants encoding different isoforms have been found for this gene.

# RBM23 Antibody (Center) Blocking Peptide - References

Lamesch, P., et al. Genomics 89(3):307-315(2007)Olsen, J.V., et al. Cell 127(3):635-648(2006)Olsen, J.V., et al. Cell 127(3):635-648(2006)Stelzl, U., et al. Cell 122(6):957-968(2005)Dowhan, D.H., et al. Mol. Cell 17(3):429-439(2005)