

HLA-C Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP17872c**Specification**

HLA-C Antibody (Center) Blocking Peptide - Product InformationPrimary Accession [P10321](#)**HLA-C Antibody (Center) Blocking Peptide - Additional Information****Gene ID** 3107**Other Names**

HLA class I histocompatibility antigen, Cw-7 alpha chain, MHC class I antigen Cw*7, HLA-C, HLAC

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

HLA-C Antibody (Center) Blocking Peptide - Protein Information**Name** HLA-C ([HGNC:4933](#))**Synonyms** HLAC**Function**

Antigen-presenting major histocompatibility complex class I (MHCI) molecule with an important role in reproduction and antiviral immunity (PubMed:20972337, PubMed:24091323, PubMed:20439706, PubMed:11172028, PubMed:20104487, PubMed:28649982, PubMed:29312307). In complex with B2M/beta 2 microglobulin displays a restricted repertoire of self and viral peptides and acts as a dominant ligand for inhibitory and activating killer immunoglobulin receptors (KIRs) expressed on NK cells (PubMed:16141329). In an allogeneic setting, such as during pregnancy, mediates interaction of extravillous trophoblasts with KIR on uterine NK cells and regulate trophoblast invasion necessary for placentation and overall fetal growth (PubMed:20972337, PubMed:<a

[24091323](http://www.uniprot.org/citations/24091323)). During viral infection, may present viral peptides with low affinity for KIRs, impeding KIR-mediated inhibition through peptide antagonism and favoring lysis of infected cells (PubMed: [20439706](http://www.uniprot.org/citations/20439706)). Presents a restricted repertoire of viral peptides on antigen-presenting cells for recognition by alpha-beta T cell receptor (TCR) on HLA-C-restricted CD8-positive T cells, guiding antigen-specific T cell immune response to eliminate infected cells, particularly in chronic viral infection settings such as HIV-1 or CMV infection (PubMed: [11172028](http://www.uniprot.org/citations/11172028) [20104487](http://www.uniprot.org/citations/20104487) [28649982](http://www.uniprot.org/citations/28649982)). Both the peptide and the MHC molecule are recognized by TCR, the peptide is responsible for the fine specificity of antigen recognition and MHC residues account for the MHC restriction of T cells (By similarity). Typically presents intracellular peptide antigens of 9 amino acids that arise from cytosolic proteolysis via proteasome. Can bind different peptides containing allele-specific binding motifs, which are mainly defined by anchor residues at position 2 and 9. Preferentially displays peptides having a restricted repertoire of hydrophobic or aromatic amino acids (Phe, Ile, Leu, Met, Val and Tyr) at the C-terminal anchor (PubMed: [8265661](http://www.uniprot.org/citations/8265661) [25311805](http://www.uniprot.org/citations/25311805)).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Endoplasmic reticulum membrane; Single-pass membrane protein

Tissue Location

Ubiquitous. Highly expressed in fetal extravillous trophoblasts in the decidua basalis (at protein level)

HLA-C Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

HLA-C Antibody (Center) Blocking Peptide - Images

HLA-C Antibody (Center) Blocking Peptide - Background

HLA-C belongs to the HLA class I heavy chain paralogues. This class I molecule is a heterodimer consisting of a heavy chain and a light chain (beta-2 microglobulin). The heavy chain is anchored in the membrane. Class I molecules play a central role in the immune system by presenting peptides derived from endoplasmic reticulum lumen. They are expressed in nearly all cells. The heavy chain is approximately 45 kDa and its gene contains 8 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the alpha1 and alpha2 domain, which both bind the peptide, exon 4 encodes the alpha3 domain, exon 5 encodes the transmembrane region, and exons 6 and 7 encode the cytoplasmic tail. Polymorphisms within exon 2 and exon 3 are responsible for the peptide binding specificity of each class one molecule. Typing for these polymorphisms is routinely done for bone marrow and kidney transplantation. Over one hundred HLA-C alleles have been described

HLA-C Antibody (Center) Blocking Peptide - References

Martin, M.P., et al. Immunogenetics 62 (11-12), 761-765 (2010) ; Strange, A., et al. Nat. Genet. 42(11):985-990(2010) Noble, J.A., et al. Diabetes 59(11):2972-2979(2010) Honeyborne, I., et al. J. Virol. 84(21):11279-11288(2010) Healy, B.C., et al. Neurology 75(7):634-640(2010)