

FNTA Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP18056a**Specification**

FNTA Antibody (N-term) Blocking Peptide - Product InformationPrimary Accession [P49354](#)**FNTA Antibody (N-term) Blocking Peptide - Additional Information****Gene ID** 2339**Other Names**

Protein farnesyltransferase/geranylgeranyltransferase type-1 subunit alpha, CAAX
farnesyltransferase subunit alpha, FTase-alpha, Ras proteins prenyltransferase subunit alpha, Type
I protein geranyl-geranyltransferase subunit alpha, GGTase-I-alpha, FNTA

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using
the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

FNTA Antibody (N-term) Blocking Peptide - Protein Information**Name** FNTA**Function**

Essential subunit of both the farnesyltransferase and the geranylgeranyltransferase complex.
Contributes to the transfer of a farnesyl or geranylgeranyl moiety from farnesyl or geranylgeranyl
diphosphate to a cysteine at the fourth position from the C-terminus of several proteins having the
C-terminal sequence Cys-aliphatic- aliphatic-X. May positively regulate neuromuscular junction
development downstream of MUSK via its function in RAC1 prenylation and activation.

FNTA Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

FNTA Antibody (N-term) Blocking Peptide - Images**FNTA Antibody (N-term) Blocking Peptide - Background**

Prenyltransferases can attach either a farnesyl group or geranylgeranyl group in thioether linkage to the cysteine residue of proteins with a C-terminal CAAX box. CAAXgeranylgeranyltransferase and CAAX farnesyltransferase are heterodimers that share the same alpha subunit but have different beta subunits. This gene encodes the alpha subunit of these transferases. Alternative splicing results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 11 and 13.

FNTA Antibody (N-term) Blocking Peptide - References

Lipkin, S.M., et al. Cancer Prev Res (Phila Pa) 3(5):597-603(2010) Fontaine-Bisson, B., et al. J. Mol. Med. 88(2):193-201(2010) Zhou, J., et al. J. Biol. Chem. 284(15):9648-9655(2009) Veluthakal, R., et al. Diabetes 56(1):204-210(2007) Armstrong, S.A., et al. J. Biol. Chem. 270(14):7864-7868(1995)