10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

UGT2B7 Antibody (N-term) Blocking Peptide

Synthetic peptide
Catalog # BP18210a

Specification

UGT2B7 Antibody (N-term) Blocking Peptide - Product Information

Primary Accession P16662

UGT2B7 Antibody (N-term) Blocking Peptide - Additional Information

Gene ID 7364

Other Names
UDP-glucuronosyltransferase 2B7, UDPGT 2B7, 4-catechol estrogen-specific UDPGT, UDP-glucuronosyltransferase 2B9, UDPGT 2B9, UDPGTh-2, UGT2B7, UGTB2B9

Format
Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage
Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions
This product is for research use only. Not for use in diagnostic or therapeutic procedures.

UGT2B7 Antibody (N-term) Blocking Peptide - Protein Information

Name UGT2B7 (HGNC:12554)

Synonyms UGTB2B9

Function
UDP-glucuronosyltransferase (UGT) that catalyzes phase II biotransformation reactions in which lipophilic substrates are conjugated with glucuronic acid to increase the metabolite's water solubility, thereby facilitating excretion into either the urine or bile (PubMed:10702251, PubMed:15472229, PubMed:15470161, PubMed:18674515, PubMed:18719240, PubMed:19022937, PubMed:23288867, PubMed:23756265, PubMed:26220143, PubMed:17442341). Essential for the elimination and detoxification of drugs, xenobiotics and endogenous compounds (PubMed:15470161, PubMed:<a

href="http://www.uniprot.org/citations/18674515" target="_blank">18674515, PubMed:23756265). Catalyzes the glucuronidation of endogenous steroid hormones such as androgens (epitestosterone, androsterone) and estrogens (estradiol, epiestradiol, estriol, catechol estrogens) (PubMed:2159463, PubMed:15472229, PubMed:18719240, PubMed:19022937, PubMed:23288867, PubMed:26220143, PubMed:17442341). Also regulates the levels of retinoic acid, a major metabolite of vitamin A involved in apoptosis, cellular growth and differentiation, and embryonic development (PubMed:10702251). Contributes to bile acid (BA) detoxification by catalyzing the glucuronidation of BA substrates, which are natural detergents for dietary lipids absorption (PubMed:23756265). Involved in the glucuronidation of the AGTR1 angiotensin receptor antagonist losartan, caderastan and zolarsatan, drugs which can inhibit the effect of angiotensin II (PubMed:18674515). Also metabolizes mycophenolate, an immunosuppressive agent (PubMed:15470161).

Cellular Location
Endoplasmic reticulum membrane; Single-pass membrane protein

UGT2B7 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Blocking Peptides

UGT2B7 Antibody (N-term) Blocking Peptide - Images

UGT2B7 Antibody (N-term) Blocking Peptide - Background

The UGTs (EC 2.4.1.17) serve a major role in theconjugation and subsequent elimination of potentially toxicxenobiotics and endogenous compounds. UGT2B7 has unique specificityfor 3,4-catechol estrogens and estriol, suggesting that it may playan important role in regulating the level and activity of thesepotent estrogen metabolites. Its subcellular location is themicrosome.

UGT2B7 Antibody (N-term) Blocking Peptide - References

Joy, M.S., et al. Eur. J. Clin. Pharmacol. 66(11):1119-1130(2010)Canzian, F., et al. Hum. Mol. Genet. 19(19):3873-3884(2010)Hu, M., et al. Pharmacogenet. Genomics 20(10):634-637(2010)Woillard, J.B., et al. Br J Clin Pharmacol 69(6):675-683(2010)Hwang, M.S., et al. Drug Metab. Pharmacokinet. 25(4):398-402(2010)