

EDA2R Antibody (C-term) Blocking Peptide

Synthetic peptide Catalog # BP18402b

Specification

EDA2R Antibody (C-term) Blocking Peptide - Product Information

Primary Accession

Q9HAV5

EDA2R Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 60401

Other Names

Tumor necrosis factor receptor superfamily member 27, X-linked ectodysplasin-A2 receptor, EDA-A2 receptor, EDA2R, TNFRSF27, XEDAR

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

EDA2R Antibody (C-term) Blocking Peptide - Protein Information

Name EDA2R

Synonyms TNFRSF27, XEDAR

Function

Receptor for EDA isoform A2, but not for EDA isoform A1. Mediates the activation of the NF-kappa-B and JNK pathways. Activation seems to be mediated by binding to TRAF3 and TRAF6.

Cellular Location

Membrane; Single-pass type III membrane protein.

EDA2R Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

EDA2R Antibody (C-term) Blocking Peptide - Images



Tel: 858.875.1900 Fax: 858.875.1999

EDA2R Antibody (C-term) Blocking Peptide - Background

EDA-A1 and EDA-A2 are two isoforms of ectodysplasin that are encoded by the anhidrotic ectodermal dysplasia (EDA) gene. Mutations in EDA give rise to a clinical syndrome characterized byloss of hair, sweat glands, and teeth. The protein encoded by thisgene specifically binds to EDA-A2 isoform. This protein is a typeIII transmembrane protein of the TNFR (tumor necrosis factorreceptor) superfamily, and contains 3 cysteine-rich repeats and asingle transmembrane domain but lacks an N-terminal signal peptide. Multiple alternatively spliced transcript variants have been foundfor this gene, but some variants lack sufficient support. [providedby RefSeq].

EDA2R Antibody (C-term) Blocking Peptide - References

Brockschmidt, F.F., et al. Br. J. Dermatol. 162(4):899-903(2010)Tanikawa, C., et al. Oncogene 28(34):3081-3092(2009)Fujimoto, A., et al. J. Hum. Genet. 54(8):461-465(2009)Hillmer, A.M., et al. Hum. Genet. 126(2):255-264(2009)Richards, J.B., et al. Nat. Genet. 40(11):1282-1284(2008)