

OR10T2 Antibody (C-term) Blocking Peptide

Synthetic peptide Catalog # BP18928b

Specification

OR10T2 Antibody (C-term) Blocking Peptide - Product Information

Primary Accession

Q8NGX3

OR10T2 Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 128360

Other Names

Olfactory receptor 10T2, Olfactory receptor OR1-3, OR10T2

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

OR10T2 Antibody (C-term) Blocking Peptide - Protein Information

Name OR10T2

Function

Odorant receptor.

Cellular Location

Cell membrane; Multi-pass membrane protein.

OR10T2 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

OR10T2 Antibody (C-term) Blocking Peptide - Images

OR10T2 Antibody (C-term) Blocking Peptide - Background

Olfactory receptors interact with odorant molecules in thenose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a largefamily of





Tel: 858.875.1900 Fax: 858.875.1999

G-protein-coupled receptors (GPCR) arising from singlecoding-exon genes. Olfactory receptors share a 7-transmembranedomain structure with many neurotransmitter and hormone receptorsand are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to theolfactory receptor genes and proteins for this organism isindependent of other organisms.

OR10T2 Antibody (C-term) Blocking Peptide - References

Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004)