

**PPP2R1A Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP1943b****Specification**

---

**PPP2R1A Antibody (C-term) Blocking Peptide - Product Information**

Primary Accession [P30153](#)  
Other Accession [Q96DH3](#)

**PPP2R1A Antibody (C-term) Blocking Peptide - Additional Information**

**Gene ID** 5518

**Other Names**

Serine/threonine-protein phosphatase 2A 65 kDa regulatory subunit A alpha isoform, Medium tumor antigen-associated 61 kDa protein, PP2A subunit A isoform PR65-alpha, PP2A subunit A isoform R1-alpha, PPP2R1A

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP1943b](/product/products/AP1943b) was selected from the C-term region of human PPP2R1A. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**PPP2R1A Antibody (C-term) Blocking Peptide - Protein Information**

**Name** PPP2R1A

**Function**

The PR65 subunit of protein phosphatase 2A serves as a scaffolding molecule to coordinate the assembly of the catalytic subunit and a variable regulatory B subunit. Upon interaction with GNA12 promotes dephosphorylation of microtubule associated protein TAU/MAPT (PubMed: [15525651](http://www.uniprot.org/citations/15525651)). Required for proper chromosome segregation and for centromeric localization of SGO1 in mitosis (PubMed: [16580887](http://www.uniprot.org/citations/16580887)). Together with RACK1 adapter, mediates dephosphorylation of AKT1 at 'Ser-473', preventing AKT1 activation and AKT-mTOR signaling pathway (By similarity). Dephosphorylation of AKT1 is essential for regulatory T-cells (Treg) homeostasis and stability (By similarity).

**Cellular Location**

Cytoplasm {ECO:0000250|UniProtKB:Q32PI5}. Nucleus. Chromosome, centromere. Lateral cell membrane. Cell projection, dendrite. Note=Centromeric localization requires the presence of BUB1.

**PPP2R1A Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**PPP2R1A Antibody (C-term) Blocking Peptide - Images****PPP2R1A Antibody (C-term) Blocking Peptide - Background**

PPP2R1A encodes a constant regulatory subunit of protein phosphatase 2. Protein phosphatase 2 is one of the four major Ser/Thr phosphatases, and it is implicated in the negative control of cell growth and division. It consists of a common heteromeric core enzyme, which is composed of a catalytic subunit and a constant regulatory subunit, that associates with a variety of regulatory subunits. The constant regulatory subunit A serves as a scaffolding molecule to coordinate the assembly of the catalytic subunit and a variable regulatory B subunit. This gene encodes an alpha isoform of the constant regulatory subunit A.