

USH1C Antibody(N-term) Blocking peptide
Synthetic peptide
Catalog # BP19566a**Specification**

USH1C Antibody(N-term) Blocking peptide - Product InformationPrimary Accession [Q9Y6N9](#)**USH1C Antibody(N-term) Blocking peptide - Additional Information****Gene ID** 10083**Other Names**

Harmonin, Antigen NY-CO-38/NY-CO-37, Autoimmune enteropathy-related antigen AIE-75, Protein PDZ-73, Renal carcinoma antigen NY-REN-3, Usher syndrome type-1C protein, USH1C, AIE75

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

USH1C Antibody(N-term) Blocking peptide - Protein Information**Name** USH1C**Synonyms** AIE75**Function**

Anchoring/scaffolding protein that is a part of the functional network formed by USH1C, USH1G, CDH23 and MYO7A that mediates mechanotransduction in cochlear hair cells. Required for normal development and maintenance of cochlear hair cell bundles (By similarity). As part of the intermicrovillar adhesion complex/IMAC plays a role in brush border differentiation, controlling microvilli organization and length. Probably plays a central regulatory role in the assembly of the complex, recruiting CDHR2, CDHR5 and MYO7B to the microvilli tips (PubMed:24725409, PubMed:26812018).

Cellular Location

Cytoplasm, cytosol. Cytoplasm, cytoskeleton. Cell projection, microvillus Note=Colocalizes with F-actin (By similarity). Detected at the tip of cochlear hair cell stereocilia (By similarity). Enriched in microvilli of the intestinal brush border (PubMed:24725409, PubMed:32209652) {ECO:0000250|UniProtKB:Q9ES64, ECO:0000269|PubMed:24725409, ECO:0000269|PubMed:32209652}

Tissue Location

Expressed in small intestine, colon, kidney, eye and weakly in pancreas. Expressed also in vestibule of the inner ear

USH1C Antibody(N-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

USH1C Antibody(N-term) Blocking peptide - Images**USH1C Antibody(N-term) Blocking peptide - Background**

This gene encodes a scaffold protein that functions in the assembly of Usher protein complexes. The protein contains PDZ domains, a coiled-coil region with a bipartite nuclear localization signal and a PEST degradation sequence. Defects in this gene are the cause of Usher syndrome type 1C and non-syndromic sensorineural deafness autosomal recessive type 18. Multiple transcript variants encoding different isoforms have been found for this gene.

USH1C Antibody(N-term) Blocking peptide - References

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) ; Yan, J., et al. Proc. Natl. Acad. Sci. U.S.A. 107(9):4040-4045(2010) ; Jaijo, T., et al. Invest. Ophthalmol. Vis. Sci. 51(3):1311-1317(2010) ; Pan, L., et al. Proc. Natl. Acad. Sci. U.S.A. 106(14):5575-5580(2009) ; Baux, D., et al. Hum. Mutat. 29 (8), E76-E87 (2008) :