

MEFV Antibody(Center) Blocking peptide
Synthetic peptide
Catalog # BP19672c**Specification**

MEFV Antibody(Center) Blocking peptide - Product InformationPrimary Accession [O15553](#)**MEFV Antibody(Center) Blocking peptide - Additional Information****Gene ID** 4210**Other Names**

Pyrin, Marenosttrin, MEFV, MEF

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

MEFV Antibody(Center) Blocking peptide - Protein Information**Name** MEFV {ECO:0000303|PubMed:11115844, ECO:0000312|HGNC:HGNC:6998}**Function**

Involved in the regulation of innate immunity and the inflammatory response in response to IFNG/IFN-gamma (PubMed:10807793, PubMed:11468188, PubMed:17964261, PubMed:18577712, PubMed:19109554, PubMed:19584923, PubMed:16037825, PubMed:27030597, PubMed:28835462, PubMed:16785446, PubMed:17431422, PubMed:26347139). Organizes autophagic machinery by serving as a platform for the assembly of ULK1, Beclin 1/BECN1, ATG16L1, and ATG8 family members and recognizes specific autophagy targets, thus coordinating target recognition with assembly of the autophagic apparatus and initiation of autophagy (PubMed:16785446, PubMed:<a

[17431422](http://www.uniprot.org/citations/17431422) PubMed: [26347139](http://www.uniprot.org/citations/26347139)). Acts as an autophagy receptor for the degradation of several inflammasome components, including CASP1, NLRP1 and NLRP3, hence preventing excessive IL1B- and IL18-mediated inflammation (PubMed: [16785446](http://www.uniprot.org/citations/16785446) PubMed: [17431422](http://www.uniprot.org/citations/17431422) PubMed: [26347139](http://www.uniprot.org/citations/26347139)). However, it can also have a positive effect in the inflammatory pathway, acting as an innate immune sensor that triggers PYCARD/ASC specks formation, caspase-1 activation, and IL1B and IL18 production (PubMed: [16037825](http://www.uniprot.org/citations/16037825) PubMed: [27030597](http://www.uniprot.org/citations/27030597) PubMed: [28835462](http://www.uniprot.org/citations/28835462)). Together with AIM2, also acts as a mediator of pyroptosis, necroptosis and apoptosis (PANoptosis), an integral part of host defense against pathogens, in response to bacterial infection (By similarity). It is required for PSTPIP1-induced PYCARD/ASC oligomerization and inflammasome formation (PubMed: [10807793](http://www.uniprot.org/citations/10807793) target="_blank">10807793 PubMed: [11468188](http://www.uniprot.org/citations/11468188) target="_blank">11468188 PubMed: [17964261](http://www.uniprot.org/citations/17964261) target="_blank">17964261 PubMed: [18577712](http://www.uniprot.org/citations/18577712) target="_blank">18577712 PubMed: [19109554](http://www.uniprot.org/citations/19109554) target="_blank">19109554 PubMed: [19584923](http://www.uniprot.org/citations/19584923) target="_blank">19584923). Recruits PSTPIP1 to inflammasomes, and is required for PSTPIP1 oligomerization (PubMed: [10807793](http://www.uniprot.org/citations/10807793) target="_blank">10807793 PubMed: [11468188](http://www.uniprot.org/citations/11468188) target="_blank">11468188 PubMed: [17964261](http://www.uniprot.org/citations/17964261) target="_blank">17964261 PubMed: [18577712](http://www.uniprot.org/citations/18577712) target="_blank">18577712 PubMed: [19109554](http://www.uniprot.org/citations/19109554) target="_blank">19109554 PubMed: [19584923](http://www.uniprot.org/citations/19584923) target="_blank">19584923).

Cellular Location

[Isoform 1]: Cytoplasm, cytoskeleton. Cell projection, ruffle. Cell projection, lamellipodium. Nucleus. Cytoplasm. Cytoplasmic vesicle, autophagosome. Note=Associated with microtubules and with the filamentous actin of perinuclear filaments and peripheral lamellar ruffles (PubMed:11468188). In pre- apoptotic cells, colocalizes with PYCARD/ASC in large specks (inflammasomes) (PubMed:11468188). In migrating monocytes, strongly polarized at the leading edge of the cell where it colocalizes with polymerizing actin and PYCARD/ASC (PubMed:11468188)

Tissue Location

Expressed in peripheral blood leukocytes, particularly in mature granulocytes and to a lesser extent in monocytes but not in lymphocytes. Detected in spleen, lung and muscle, probably as a result of leukocyte infiltration in these tissues. Not expressed in thymus, prostate, testis, ovary, small intestine, colon, heart, brain, placenta, liver, kidney, pancreas. Expression detected in several myeloid leukemic, colon cancer, and prostate cancer cell lines

MEFV Antibody(Center) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

MEFV Antibody(Center) Blocking peptide - Images

MEFV Antibody(Center) Blocking peptide - Background

This gene encodes a protein, also known as pyrin or marenostrin, that is an important modulator of innate immunity. Mutations in this gene are associated with Mediterranean fever, a hereditary

periodic fever syndrome.

MEFV Antibody(Center) Blocking peptide - References

Cosan, F., et al. Arthritis Rheum. 62(11):3232-3236(2010)Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)He, X., et al. Pediatr. Nephrol. 25(10):2077-2082(2010)Simsek, I., et al. Clin. Rheumatol. (2010) In press :