

NCOA3 Blocking Peptide (N-term)

Synthetic peptide

Catalog # BP19843a

Specification

NCOA3 Blocking Peptide (N-term) - Product Information

Primary Accession

[O9Y6Q9](#)

Other Accession

[O9EPU2](#), [NP_001167559.1](#)**NCOA3 Blocking Peptide (N-term) - Additional Information****Gene ID** 8202**Other Names**

Nuclear receptor coactivator 3, NCoA-3, ACTR, Amplified in breast cancer 1 protein, AIB-1, CBP-interacting protein, pCIP, Class E basic helix-loop-helix protein 42, bHLHe42, Receptor-associated coactivator 3, RAC-3, Steroid receptor coactivator protein 3, SRC-3, Thyroid hormone receptor activator molecule 1, TRAM-1, NCOA3, AIB1, BHLHE42, RAC3, TRAM1

Target/Specificity

The synthetic peptide sequence is selected from aa 359-372 of HUMAN NCOA3

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

NCOA3 Blocking Peptide (N-term) - Protein Information**Name** NCOA3**Synonyms** AIB1, BHLHE42, RAC3, TRAM1**Function**

Nuclear receptor coactivator that directly binds nuclear receptors and stimulates the transcriptional activities in a hormone- dependent fashion. Plays a central role in creating a multisubunit coactivator complex, which probably acts via remodeling of chromatin. Involved in the coactivation of different nuclear receptors, such as for steroids (GR and ER), retinoids (RARs and RXRs), thyroid hormone (TRs), vitamin D3 (VDR) and prostanoids (PPARs). Displays histone acetyltransferase activity. Also involved in the coactivation of the NF-kappa-B pathway via its interaction with the NFKB1 subunit.

Cellular Location

Cytoplasm. Nucleus. Note=Mainly cytoplasmic and weakly nuclear. Upon TNF activation and subsequent phosphorylation, it translocates from the cytoplasm to the nucleus

Tissue Location

Widely expressed. High expression in heart, skeletal muscle, pancreas and placenta. Low expression in brain, and very low in lung, liver and kidney

NCOA3 Blocking Peptide (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

NCOA3 Blocking Peptide (N-term) - Images**NCOA3 Blocking Peptide (N-term) - Background**

The protein encoded by this gene is a nuclear receptor coactivator that interacts with nuclear hormone receptors to enhance their transcriptional activator functions. The encoded protein has histone acetyltransferase activity and recruits p300/CBP-associated factor and CREB binding protein as part of a multisubunit coactivation complex. This protein is initially found in the cytoplasm but is translocated into the nucleus upon phosphorylation. Several transcript variants encoding different isoforms have been found for this gene. In addition, a polymorphic repeat region is found in the C-terminus of the encoded protein.

NCOA3 Blocking Peptide (N-term) - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Cai, D., et al. Cancer Res. 70(16):6477-6485(2010)
Kovanen, L., et al. Alcohol Alcohol. 45(4):303-311(2010)
Karmakar, S., et al. Mol. Endocrinol. 24(6):1187-1202(2010)
Kleibl, Z., et al. J. Cancer Res. Clin. Oncol. (2010) In press :