

**UNC5C Blocking Peptide (N-term)**  
**Synthetic peptide**  
**Catalog # BP19910a****Specification**

---

**UNC5C Blocking Peptide (N-term) - Product Information**

Primary Accession [O95185](#)  
Other Accession [NP\\_003719.2](#)

**UNC5C Blocking Peptide (N-term) - Additional Information**

**Gene ID** 8633

**Other Names**

Netrin receptor UNC5C, Protein unc-5 homolog 3, Protein unc-5 homolog C, UNC5C, UNC5H3

**Target/Specificity**

The synthetic peptide sequence is selected from aa 103-117 of HUMAN UNC5C

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**UNC5C Blocking Peptide (N-term) - Protein Information**

**Name** UNC5C

**Synonyms** UNC5H3

**Function**

Receptor for netrin required for axon guidance (By similarity). Mediates axon repulsion of neuronal growth cones in the developing nervous system upon ligand binding (By similarity). NTN1/Netrin-1 binding might cause dissociation of UNC5C from polymerized TUBB3 in microtubules and thereby lead to increased microtubule dynamics and axon repulsion (PubMed:<a href="http://www.uniprot.org/citations/28483977" target="\_blank">28483977</a>). Axon repulsion in growth cones may also be caused by its association with DCC that may trigger signaling for repulsion (By similarity). Might also collaborate with DSCAM in NTN1-mediated axon repulsion independently of DCC (By similarity). Also involved in corticospinal tract axon guidance independently of DCC (By similarity). Involved in dorsal root ganglion axon projection towards the spinal cord (PubMed:<a href="http://www.uniprot.org/citations/28483977" target="\_blank">28483977</a>). It also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand (By similarity).

**Cellular Location**

Cell membrane; Single-pass type I membrane protein. Cell surface. Synapse, synaptosome {ECO:0000250|UniProtKB:Q761X5}. Cell projection, axon {ECO:0000250|UniProtKB:O08747}. Cell projection, dendrite {ECO:0000250|UniProtKB:O08747}. Cell projection, growth cone {ECO:0000250|UniProtKB:O08747}. Cell projection, lamellipodium {ECO:0000250|UniProtKB:O08747}. Cell projection, filopodium {ECO:0000250|UniProtKB:O08747}

**Tissue Location**

Mainly expressed in brain (PubMed:9782087). Expressed in temporal lobe cortical neurons and in neurons of the hippocampal pyramidal layer (PubMed:25419706). Also expressed in kidney (PubMed:9782087). Not expressed in developing or adult lung (PubMed:9782087).

**UNC5C Blocking Peptide (N-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**UNC5C Blocking Peptide (N-term) - Images****UNC5C Blocking Peptide (N-term) - Background**

Receptor for netrin required for axon guidance. Mediates axon repulsion of neuronal growth cones in the developing nervous system upon ligand binding. Axon repulsion in growth cones may be caused by its association with DCC that may trigger signaling for repulsion. Also involved in corticospinal tract axon guidances independently of DCC. It also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand (By similarity).