

# Neurogenin3 Antibody (C-term) Blocking Peptide

Synthetic peptide Catalog # BP2024b

## **Specification**

## Neurogenin3 Antibody (C-term) Blocking Peptide - Product Information

**Primary Accession** 

**09Y4Z2** 

# Neurogenin3 Antibody (C-term) Blocking Peptide - Additional Information

**Gene ID 50674** 

#### **Other Names**

Neurogenin-3, NGN-3, Class A basic helix-loop-helix protein 7, bHLHa7, Protein atonal homolog 5, NEUROG3, ATOH5, BHLHA7, NGN3

## **Target/Specificity**

The synthetic peptide sequence used to generate the antibody <a href=/product/products/AP2024b>AP2024b</a> was selected from the C-term region of human Neurogenin3 . A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### Neurogenin3 Antibody (C-term) Blocking Peptide - Protein Information

Name NEUROG3

Synonyms ATOH5, BHLHA7, NGN3

### **Function**

Acts as a transcriptional regulator. Together with NKX2-2, initiates transcriptional activation of NEUROD1. Involved in neurogenesis. Also required for the specification of a common precursor of the 4 pancreatic endocrine cell types (By similarity).

### **Cellular Location**

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00981}.



## Neurogenin3 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

# • Blocking Peptides

Neurogenin3 Antibody (C-term) Blocking Peptide - Images

# Neurogenin3 Antibody (C-term) Blocking Peptide - Background

Neurogenin-3 belongs to a family of basic helix-loop-helix transcription factors involved in the determination of neural precursor cells in the neuroectoderm. Neurogenin-3 also required for the specification of a common precursor of the 4 pancreatic endocrine cell types. Defects in NEUROG3 are the cause of congenital malabsorptive diarrhea 4 (DIAR4). DIAR4 is an autosomal recessive disorder characterized by generalized malabsorption and a paucity of enteroendocrine cells.

## Neurogenin3 Antibody (C-term) Blocking Peptide - References

Heremans, Y., et al., J. Cell Biol. 159(2):303-312 (2002). Sommer, L., et al., Mol. Cell. Neurosci. 8(4):221-241 (1996).