

**GFI1B Blocking Peptide (C-term)**

Synthetic peptide

Catalog # BP20437b

**Specification**

---

**GFI1B Blocking Peptide (C-term) - Product Information**

Primary Accession

[Q5VTD9](#)

Other Accession

[O70237](#)**GFI1B Blocking Peptide (C-term) - Additional Information**

Gene ID 8328

**Other Names**

Zinc finger protein Gfi-1b, Growth factor independent protein 1B, Potential regulator of CDKN1A translocated in CML, GFI1B

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**GFI1B Blocking Peptide (C-term) - Protein Information**

Name GFI1B

**Function**

Essential proto-oncogenic transcriptional regulator necessary for development and differentiation of erythroid and megakaryocytic lineages. Component of a RCOR-GFI-KDM1A-HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development and controls hematopoietic differentiation. Transcriptional repressor or activator depending on both promoter and cell type context; represses promoter activity of SOCS1 and SOCS3 and thus, may regulate cytokine signaling pathways. Cooperates with GATA1 to repress target gene transcription, such as the apoptosis regulator BCL2L1; GFI1B silencing in leukemic cell lines markedly increase apoptosis rate. Inhibits down-regulation of MYC and MYB as well as the cyclin-dependent kinase inhibitor CDKN1A/P21WAF1 in IL6-treated myelomonocytic cells. Represses expression of GATA3 in T-cell lymphomas and inhibits GATA1-mediated transcription; as GATA1 also mediates erythroid GFI1B transcription, both GATA1 and GFI1B participate in a feedback regulatory pathway controlling the expression of GFI1B gene in erythroid cells. Suppresses GATA1-mediated stimulation of GFI1B promoter through protein interaction. Binds to gamma-satellite DNA and to its own promoter, auto-repressing its own expression. Alters histone methylation by recruiting histone methyltransferase to target genes promoters. Plays a role in heterochromatin formation.

**Cellular Location**

Nucleus.

**Tissue Location**

Expressed in bone marrow and fetal liver, but also detectable in fetal spleen, fetal thymus, and testes. Detected in hematopoietic stem cells, erythroblasts, and megakaryocytes Overexpressed in bone marrow of patients with erythroleukemia and megakaryocytic leukemia as well as in their corresponding leukemic cell lines, and markedly repressed in severe aplastic anemia (SAA)

**GFI1B Blocking Peptide (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**GFI1B Blocking Peptide (C-term) - Images****GFI1B Blocking Peptide (C-term) - Background**

Essential proto-oncogenic transcriptional regulator necessary for development and differentiation of erythroid and megakaryocytic lineages. Component of a RCOR-GFI-KDM1A-HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development and controls hematopoietic differentiation. Transcriptional repressor or activator depending on both promoter and cell type context; represses promoter activity of SOCS1 and SOCS3 and thus, may regulate cytokine signaling pathways. Cooperates with GATA1 to repress target gene transcription, such as the apoptosis regulator BCL2L1; GFI1B silencing in leukemic cell lines markedly increase apoptosis rate. Inhibits down-regulation of MYC and MYB as well as the cyclin-dependent kinase inhibitor CDKN1A/P21WAF1 in IL6-treated myelomonocytic cells. Represses expression of GATA3 in T-cell lymphomas and inhibits GATA1-mediated transcription; as GATA1 also mediates erythroid GFI1B transcription, both GATA1 and GFI1B participate in a feedback regulatory pathway controlling the expression of GFI1B gene in erythroid cells. Suppresses GATA1-mediated stimulation of GFI1B promoter through protein interaction. Binds to gamma-satellite DNA and to its own promoter, auto-repressing its own expression. Alters histone methylation by recruiting histone methyltransferase to target genes promoters. Plays a role in heterochromatin formation.

**GFI1B Blocking Peptide (C-term) - References**

Roedel B., et al. Genomics 54:580-582(1998).  
Huang D.Y., et al. Nucleic Acids Res. 32:3935-3946(2004).  
Halleck A., et al. Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.  
Humphray S.J., et al. Nature 429:369-374(2004).  
Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.