

OCT4 (OCT3) Antibody (C-term) Blocking peptide
Synthetic peptide
Catalog # BP2046b**Specification**

OCT4 (OCT3) Antibody (C-term) Blocking peptide - Product Information

Primary Accession [O01860](#)
Other Accession [NP_002692](#)

OCT4 (OCT3) Antibody (C-term) Blocking peptide - Additional Information

Gene ID 5460

Other Names

POU domain, class 5, transcription factor 1, Octamer-binding protein 3, Oct-3, Octamer-binding protein 4, Oct-4, Octamer-binding transcription factor 3, OTF-3, POU5F1, OCT3, OCT4, OTF3

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP2046b](/product/products/AP2046b) was selected from the OCT4 region of human OCT3 (OCT4). A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

OCT4 (OCT3) Antibody (C-term) Blocking peptide - Protein Information

Name POU5F1

Synonyms OCT3, OCT4, OTF3

Function

Transcription factor that binds to the octamer motif (5'- ATTTGCAT-3'). Forms a trimeric complex with SOX2 or SOX15 on DNA and controls the expression of a number of genes involved in embryonic development such as YES1, FGF4, UTF1 and ZFP206. Critical for early embryogenesis and for embryonic stem cell pluripotency.

Cellular Location

Cytoplasm. Nucleus. Note=Expressed in a diffuse and slightly punctuate pattern. Colocalizes with MAPK8 and MAPK9 in the nucleus. {ECO:0000250|UniProtKB:P20263,

ECO:0000269|PubMed:18191611, ECO:0000269|PubMed:19274063,
ECO:0000269|PubMed:23024368}

Tissue Location

Expressed in developing brain. Highest levels found in specific cell layers of the cortex, the olfactory bulb, the hippocampus and the cerebellum. Low levels of expression in adult tissues.

OCT4 (OCT3) Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

OCT4 (OCT3) Antibody (C-term) Blocking peptide - Images**OCT4 (OCT3) Antibody (C-term) Blocking peptide - Background**

Transcription factors containing the POU homeodomain have been shown to be important regulators of tissue-specific gene expression in lymphoid and pituitary differentiation and in early mammalian development. Two forms of OCT3 mRNA are expressed in adult tissues as a result of alternative splicing--OCT3A and OCT3B. Reverse transcriptase PCR showed low level of expression in both OCT3A and OCT3B mRNA in all adult human tissues examined. Oct3 is present in mouse oocytes before and after fertilization. When fertilized oocytes were injected with antisense Oct3 oligonucleotides or double-stranded DNA containing the octamer motif, embryonic DNA synthesis was inhibited and the embryos were arrested at the one-cell stage.