

Mouse Hck Blocking Peptide (Center)
Synthetic peptide
Catalog # BP20682c**Specification**

Mouse Hck Blocking Peptide (Center) - Product Information

Primary Accession [P08103](#)
Other Accession [P50545](#)

Mouse Hck Blocking Peptide (Center) - Additional Information

Gene ID 15162

Other Names

Tyrosine-protein kinase HCK, B-cell/myeloid kinase, BMK, Hematopoietic cell kinase, Hemopoietic cell kinase, p56-HCK/p59-HCK, Hck

Target/Specificity

The synthetic peptide sequence is selected from aa 239-252 of HUMAN Hck

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Mouse Hck Blocking Peptide (Center) - Protein Information

Name Hck

Function

Non-receptor tyrosine-protein kinase found in hematopoietic cells that transmits signals from cell surface receptors and plays an important role in the regulation of innate immune responses, including neutrophil, monocyte, macrophage and mast cell functions, phagocytosis, cell survival and proliferation, cell adhesion and migration. Acts downstream of receptors that bind the Fc region of immunoglobulins, such as FCGR1A and FCGR2A, but also CSF3R, PLAUR, the receptors for IFNG, IL2, IL6 and IL8, and integrins, such as ITGB1 and ITGB2. During the phagocytic process, mediates mobilization of secretory lysosomes, degranulation, and activation of NADPH oxidase to bring about the respiratory burst. Plays a role in the release of inflammatory molecules. Promotes reorganization of the actin cytoskeleton and actin polymerization, formation of podosomes and cell protrusions. Inhibits TP73-mediated transcription activation and TP73-mediated apoptosis. Phosphorylates CBL in response to activation of immunoglobulin gamma Fc region receptors. Phosphorylates ADAM15, BCR, ELMO1, FCGR2A, GAB1, GAB2, RAPGEF1, STAT5B, TP73, VAV1 and WAS (By similarity).

Cellular Location

Cytoplasmic vesicle, secretory vesicle. Cytoplasm, cytosol [Isoform 2]: Cell membrane; Lipid-anchor. Membrane, caveola; Lipid-anchor. Cell junction, focal adhesion. Cytoplasm, cytoskeleton. Golgi apparatus. Cytoplasmic vesicle. Lysosome. Nucleus. Note=20% of this isoform is associated with caveolae. Localization at the cell membrane and at caveolae requires palmitoylation at Cys-3. Colocalizes with the actin cytoskeleton at focal adhesions (By similarity)

Tissue Location

Expressed predominantly in cells of the myeloid and B-lymphoid lineages

Mouse Hck Blocking Peptide (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

Mouse Hck Blocking Peptide (Center) - Images**Mouse Hck Blocking Peptide (Center) - Background**

Non-receptor tyrosine-protein kinase found in hematopoietic cells that transmits signals from cell surface receptors and plays an important role in the regulation of innate immune responses, including neutrophil, monocyte, macrophage and mast cell functions, phagocytosis, cell survival and proliferation, cell adhesion and migration. Acts downstream of receptors that bind the Fc region of immunoglobulins, such as FCGR1A and FCGR2A, but also CSF3R, PLAUR, the receptors for IFNG, IL2, IL6 and IL8, and integrins, such as ITGB1 and ITGB2. During the phagocytic process, mediates mobilization of secretory lysosomes, degranulation, and activation of NADPH oxidase to bring about the respiratory burst. Plays a role in the release of inflammatory molecules. Promotes reorganization of the actin cytoskeleton and actin polymerization, formation of podosomes and cell protrusions. Inhibits TP73-mediated transcription activation and TP73-mediated apoptosis. Phosphorylates CBL in response to activation of immunoglobulin gamma Fc region receptors. Phosphorylates ADAM15, BCR, ELMO1, FCGR2A, GAB1, GAB2, RAPGEF1, STAT5B, TP73, VAV1 and WAS (By similarity).

Mouse Hck Blocking Peptide (Center) - References

Klemsz M.J.,et al.Nucleic Acids Res. 15:9600-9600(1987).
Holtzman D.A.,et al.Proc. Natl. Acad. Sci. U.S.A. 84:8325-8329(1987).
Carninci P.,et al.Science 309:1559-1563(2005).
Lock P.,et al.Mol. Cell. Biol. 11:4363-4370(1991).
Lowell C.A.,et al.Genes Dev. 8:387-398(1994).