10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Mouse Ring1 Blocking Peptide (C-term)

Synthetic peptide
Catalog # BP20847c

Specification

Mouse Ring1 Blocking Peptide (C-term) - Product Information

Primary Accession O35730

Mouse Ring1 Blocking Peptide (C-term) - Additional Information

Gene ID 19763

Other Names
E3 ubiquitin-protein ligase RING1, 632-, Polycomb complex protein RING1, RING finger protein 1, Transcription repressor Ring1A, Ring1, Ring1A, Rnf1

Target/Specificity
The synthetic peptide sequence is selected from aa 306-320 of HUMAN Ring1

Format
Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage
Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions
This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Mouse Ring1 Blocking Peptide (C-term) - Protein Information

Name Ring1 {ECO:0000312|MGI:MGI:1101770}

Synonyms Ring1A, Rnf1

Function
Constitutes one of the E3 ubiquitin-protein ligases that mediate monoubiquitination of 'Lys-119' of histone H2A, thereby playing a central role in histone code and gene regulation. H2A 'Lys-119' ubiquitination gives a specific tag for epigenetic transcriptional repression and participates in X chromosome inactivation of female mammals. Essential component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones, rendering chromatin heritably changed in its expressibility. Compared to RNF2/RING2, it does not have the main E3 ubiquitin ligase activity on histone H2A, and it may rather act as a modulator of RNF2/RING2 activity (By similarity).

Cellular Location

Nucleus speckle.

Mouse Ring1 Blocking Peptide (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Blocking Peptides

Mouse Ring1 Blocking Peptide (C-term) - Images

Mouse Ring1 Blocking Peptide (C-term) - Background

Constitutes one of the E3 ubiquitin-protein ligases that mediate monoubiquitination of 'Lys-119' of histone H2A, thereby playing a central role in histone code and gene regulation. H2A 'Lys-119' ubiquitination gives a specific tag for epigenetic transcriptional repression and participates in X chromosome inactivation of female mammals. Essential component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones, rendering chromatin heritably changed in its expressibility. Compared to RNF2/RING2, it does not have the main E3 ubiquitin ligase activity on histone H2A, and it may rather act as a modulator of RNF2/RING2 activity (By similarity).

Mouse Ring1 Blocking Peptide (C-term) - References

Schoorlemmer J.,et al.EMBO J. 16:5930-5942(1997).
Rowen L.,et al.Submitted (OCT-1998) to the EMBL/GenBank/DDBJ databases.
Carninci P.,et al.Science 309:1559-1563(2005).
Ebert L.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
de Napoles M.,et al.Dev. Cell 7:663-676(2004).