

IHH Blocking Peptide (C-term)
Synthetic peptide
Catalog # BP20973c**Specification**

IHH Blocking Peptide (C-term) - Product InformationPrimary Accession [Q14623](#)**IHH Blocking Peptide (C-term) - Additional Information****Gene ID** 3549**Other Names**

Indian hedgehog protein, IHH, HHG-2, Indian hedgehog protein N-product, Indian hedgehog protein C-product, IHH

Target/Specificity

The synthetic peptide sequence is selected from aa 397-411 of HUMAN IHH

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

IHH Blocking Peptide (C-term) - Protein Information**Name** IHH ([HGNC:5956](#))**Function**

[Indian hedgehog protein]: The C-terminal part of the indian hedgehog protein precursor displays an autoproteolysis and a cholesterol transferase activity (By similarity). Both activities result in the cleavage of the full-length protein into two parts followed by the covalent attachment of a cholesterol moiety to the C- terminal of the newly generated N-product (By similarity). Both activities occur in the reticulum endoplasmic (By similarity). Plays a role in hedgehog paracrine signaling (PubMed:24342078). Associated with the very-low-density lipoprotein (VLDL) particles to function as a circulating morphogen for endothelial cell integrity maintenance (PubMed:20839884).

Cellular Location

[Indian hedgehog protein N-product]: Cell membrane; Lipid-anchor {ECO:0000250|UniProtKB:Q62226}. Note=The N-product remains associated with the cell surface. {ECO:0000250|UniProtKB:Q15465}

Tissue Location

Expressed in embryonic lung, and in adult kidney and liver

IHH Blocking Peptide (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

IHH Blocking Peptide (C-term) - Images**IHH Blocking Peptide (C-term) - Background**

Intercellular signal essential for a variety of patterning events during development. Binds to the patched (PTC) receptor, which functions in association with smoothened (SMO), to activate the transcription of target genes. Implicated in endochondral ossification: may regulate the balance between growth and ossification of the developing bones. Induces the expression of parathyroid hormone-related protein (PTHrP) (By similarity).

IHH Blocking Peptide (C-term) - References

Tate G.,et al.J. Biochem. Mol. Biol. Biophys. 4:27-34(2000).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Marigo V.,et al.Genomics 28:44-51(1995).
Chang D.T.,et al.Development 120:3339-3353(1994).
Pepinsky R.B.,et al.J. Biol. Chem. 273:14037-14045(1998).