

## **EWSR1 Blocking Peptide (Center)**

Synthetic peptide Catalog # BP20981a

## **Specification**

### **EWSR1 Blocking Peptide (Center) - Product Information**

Primary Accession Q01844
Other Accession Q61545

## EWSR1 Blocking Peptide (Center) - Additional Information

**Gene ID 2130** 

#### **Other Names**

RNA-binding protein EWS, EWS oncogene, Ewing sarcoma breakpoint region 1 protein, EWSR1, EWS

### **Target/Specificity**

The synthetic peptide sequence is selected from aa 328-342 of HUMAN EWSR1

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

# **EWSR1 Blocking Peptide (Center) - Protein Information**

#### Name EWSR1

## **Synonyms** EWS

## **Function**

Might normally function as a transcriptional repressor. EWS- fusion-proteins (EFPS) may play a role in the tumorigenic process. They may disturb gene expression by mimicking, or interfering with the normal function of CTD-POLII within the transcription initiation complex. They may also contribute to an aberrant activation of the fusion protein target genes.

#### **Cellular Location**

Nucleus. Cytoplasm. Cell membrane. Note=Relocates from cytoplasm to ribosomes upon PTK2B/FAK2 activation

# Tissue Location Ubiquitous.



## **EWSR1 Blocking Peptide (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

## • Blocking Peptides

**EWSR1 Blocking Peptide (Center) - Images** 

## EWSR1 Blocking Peptide (Center) - Background

Might normally function as a transcriptionnal repressor. EWS-fusion-proteins (EFPS) may play a role in the tumorigenic process. They may disturb gene expression by mimicking, or interfering with the normal function of CTD-POLII within the transcription initiation complex. They may also contribute to an aberrant activation of the fusion protein target genes.

## **EWSR1 Blocking Peptide (Center) - References**

Delattre O., et al. Nature 359:162-165(1992). Plougastel B., et al. Genomics 18:609-615(1993). Zucman-Rossi J., et al. Submitted (MAY-1998) to the EMBL/GenBank/DDBJ databases. Collins J.E., et al. Genome Biol. 5:R84.1-R84.11(2004). Ota T., et al. Nat. Genet. 36:40-45(2004).