

NDEL1 Blocking Peptide (Center)
Synthetic peptide
Catalog # BP21101a**Specification**

NDEL1 Blocking Peptide (Center) - Product Information

Primary Accession
Other Accession

[O9GZM8](#)
[Q78PB6](#), [Q46480](#), [Q9ERR1](#), [Q4R4S6](#), [Q5ZKH4](#)

NDEL1 Blocking Peptide (Center) - Additional Information

Gene ID 81565

Other Names

Nuclear distribution protein nudE-like 1, Protein Nudel, Mitosin-associated protein 1, NDEL1, EOPA, MITAP1, NUDEL

Target/Specificity

The synthetic peptide sequence is selected from aa 178-193 of HUMAN NDEL1

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

NDEL1 Blocking Peptide (Center) - Protein Information

Name NDEL1

Synonyms EOPA, MITAP1, NUDEL

Function

Required for organization of the cellular microtubule array and microtubule anchoring at the centrosome. May regulate microtubule organization at least in part by targeting the microtubule severing protein KATNA1 to the centrosome. Also positively regulates the activity of the minus-end directed microtubule motor protein dynein. May enhance dynein-mediated microtubule sliding by targeting dynein to the microtubule plus ends. Required for several dynein- and microtubule-dependent processes such as the maintenance of Golgi integrity, the centripetal motion of secretory vesicles and the coupling of the nucleus and centrosome. Also required during brain development for the migration of newly formed neurons from the ventricular/subventricular zone toward the cortical plate. Plays a role, together with DISC1, in the regulation of neurite outgrowth. Required for mitosis in some cell types but appears to be dispensable for mitosis in cortical neuronal progenitors, which instead requires NDE1. Facilitates the polymerization of

neurofilaments from the individual subunits NEFH and NEFL. Positively regulates lysosome peripheral distribution and ruffled border formation in osteoclasts (By similarity).

Cellular Location

Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Chromosome, centromere, kinetochore. Cytoplasm, cytoskeleton, spindle. Note=Localizes to the cell body of the motor neurons and colocalizes with assembled neurofilaments within axonal processes. Localizes to the microtubules of the manchette in elongated spermatids. Colocalizes with DISC1 in the perinuclear region, including the centrosome (By similarity). Localizes to the interphase centrosome and the mitotic spindle. Localizes to the kinetochore in a CENPF-dependent manner.

Tissue Location

Expressed in brain, heart, kidney, liver, lung, pancreas, placenta and skeletal muscle.

NDEL1 Blocking Peptide (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

NDEL1 Blocking Peptide (Center) - Images**NDEL1 Blocking Peptide (Center) - Background**

Required for organization of the cellular microtubule array and microtubule anchoring at the centrosome. May regulate microtubule organization at least in part by targeting the microtubule severing protein KATNA1 to the centrosome. Also positively regulates the activity of the minus-end directed microtubule motor protein dynein. May enhance dynein-mediated microtubule sliding by targeting dynein to the microtubule plus ends. Required for several dynein- and microtubule-dependent processes such as the maintenance of Golgi integrity, the centripetal motion of secretory vesicles and the coupling of the nucleus and centrosome. Also required during brain development for the migration of newly formed neurons from the ventricular/subventricular zone toward the cortical plate. Plays a role, together with DISC1, in the regulation of neurite outgrowth. Required for mitosis in some cell types but appears to be dispensable for mitosis in cortical neuronal progenitors, which instead requires NDE1. Facilitates the polymerization of neurofilaments from the individual subunits NEFH and NEFL.

NDEL1 Blocking Peptide (Center) - References

Niethammer M., et al. Neuron 28:697-711(2000).
Yan X., et al. Mol. Cell. Biol. 23:1239-1250(2003).
Hayashi M.A.F., et al. Proc. Natl. Acad. Sci. U.S.A. 102:3828-3833(2005).
Ota T., et al. Nat. Genet. 36:40-45(2004).
Bechtel S., et al. BMC Genomics 8:399-399(2007).