

**(Mouse) Med15 Blocking Peptide (C-term)**  
**Synthetic peptide**  
**Catalog # BP21208b**

## **Specification**

---

### **(Mouse) Med15 Blocking Peptide (C-term) - Product Information**

Primary Accession [Q924H2](#)

### **(Mouse) Med15 Blocking Peptide (C-term) - Additional Information**

**Gene ID** 94112

#### **Other Names**

Mediator of RNA polymerase II transcription subunit 15, Mediator complex subunit 15, Positive cofactor 2 glutamine/Q-rich-associated protein, PC2 glutamine/Q-rich-associated protein, mPcqap, Med15, Pcqap

#### **Target/Specificity**

The synthetic peptide sequence is selected from aa 712-726 of HUMAN Med15

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### **(Mouse) Med15 Blocking Peptide (C-term) - Protein Information**

**Name** Med15

**Synonyms** Pcqap

#### **Function**

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. Required for cholesterol-dependent gene regulation. Positively regulates the Nodal signaling pathway (By similarity).

#### **Cellular Location**

Cytoplasm. Nucleus.

**(Mouse) Med15 Blocking Peptide (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**(Mouse) Med15 Blocking Peptide (C-term) - Images****(Mouse) Med15 Blocking Peptide (C-term) - Background**

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. Required for cholesterol-dependent gene regulation. Positively regulates the Nodal signaling pathway (By similarity).

**(Mouse) Med15 Blocking Peptide (C-term) - References**

Berti L., et al. Genomics 74:320-332(2001).