

**(Mouse) Ehmt2 Blocking Peptide (N-term)**  
**Synthetic peptide**  
**Catalog # BP21266a**

**Specification**

---

**(Mouse) Ehmt2 Blocking Peptide (N-term) - Product Information**

Primary Accession [Q9Z148](#)

**(Mouse) Ehmt2 Blocking Peptide (N-term) - Additional Information**

**Gene ID** 110147

**Other Names**

Histone-lysine N-methyltransferase EHMT2, 211-, Euchromatic histone-lysine N-methyltransferase 2, HLA-B-associated transcript 8, Histone H3-K9 methyltransferase 3, H3-K9-HMTase 3, Protein G9a, Ehmt2, Bat8, G9a, Ng36

**Target/Specificity**

The synthetic peptide sequence is selected from aa 227-239 of HUMAN Ehmt2

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**(Mouse) Ehmt2 Blocking Peptide (N-term) - Protein Information**

**Name** Ehmt2

**Synonyms** Bat8, G9a, Ng36

**Function**

Histone methyltransferase that specifically mono- and dimethylates 'Lys-9' of histone H3 (H3K9me1 and H3K9me2, respectively) in euchromatin. H3K9me represents a specific tag for epigenetic transcriptional repression by recruiting HP1 proteins to methylated histones. Also mediates monomethylation of 'Lys-56' of histone H3 (H3K56me1) in G1 phase, leading to promote interaction between histone H3 and PCNA and regulating DNA replication. Also weakly methylates 'Lys-27' of histone H3 (H3K27me). Also required for DNA methylation, the histone methyltransferase activity is not required for DNA methylation, suggesting that these 2 activities function independently. Probably targeted to histone H3 by different DNA-binding proteins like E2F6, MGA, MAX and/or DP1. May also methylate histone H1. In addition to the histone methyltransferase activity, also methylates non-histone proteins: mediates dimethylation of 'Lys-373' of p53/TP53. Also methylates CDYL, WIZ, ACIN1, DNMT1, HDAC1, ERCC6, KLF12 and

itself. Recruited to the promoters of target genes through interaction with transcriptional repressor MSX1, leading to the inhibition of myoblast differentiation via transcriptional repression of differentiation factors (PubMed:<a href="http://www.uniprot.org/citations/22629437" target="\_blank">22629437</a>).

#### **Cellular Location**

Nucleus {ECO:0000250|UniProtKB:Q96KQ7}. Chromosome {ECO:0000250|UniProtKB:Q96KQ7}. Note=Almost excluded from nucleoli. Associates with euchromatic regions (By similarity). Does not associate with heterochromatin (By similarity) {ECO:0000250|UniProtKB:Q96KQ7}

#### **Tissue Location**

Ubiquitous..

### **(Mouse) Ehmt2 Blocking Peptide (N-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

### **(Mouse) Ehmt2 Blocking Peptide (N-term) - Images**

### **(Mouse) Ehmt2 Blocking Peptide (N-term) - Background**

Histone methyltransferase that specifically mono- and dimethylates 'Lys-9' of histone H3 (H3K9me1 and H3K9me2, respectively) in euchromatin. H3K9me represents a specific tag for epigenetic transcriptional repression by recruiting HP1 proteins to methylated histones. Also mediates monomethylation of 'Lys-56' of histone H3 (H3K56me1) in G1 phase, leading to promote interaction between histone H3 and PCNA and regulating DNA replication. Also weakly methylates 'Lys-27' of histone H3 (H3K27me). Also required for DNA methylation, the histone methyltransferase activity is not required for DNA methylation, suggesting that these 2 activities function independently. Probably targeted to histone H3 by different DNA-binding proteins like E2F6, MGA, MAX and/or DP1. May also methylate histone H1. In addition to the histone methyltransferase activity, also methylates non-histone proteins: mediates dimethylation of 'Lys-373' of p53/TP53. Also methylates CDYL, WIZ, ACIN1, DNMT1, HDAC1, ERCC6, KLF12 and itself.

### **(Mouse) Ehmt2 Blocking Peptide (N-term) - References**

Tachibana M.,et al.Genes Dev. 16:1779-1791(2002).  
Xie T.,et al.Genome Res. 13:2621-2636(2003).  
Church D.M.,et al.PLoS Biol. 7:E1000112-E1000112(2009).  
Brown S.E.,et al.Mamm. Genome 12:916-924(2001).  
Tachibana M.,et al.J. Biol. Chem. 276:25309-25317(2001).