

RBBP5 Blocking Peptide (C-term) Synthetic peptide Catalog # BP21555b

# Specification

# **RBBP5 Blocking Peptide (C-term) - Product Information**

Primary Accession

<u>Q15291</u>

# **RBBP5 Blocking Peptide (C-term) - Additional Information**

Gene ID 5929

**Other Names** Retinoblastoma-binding protein 5, RBBP-5, Retinoblastoma-binding protein RBQ-3, RBBP5, RBQ3

### **Target/Specificity**

The synthetic peptide sequence is selected from aa 505-539 of HUMAN RBBP5

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage** Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### **RBBP5 Blocking Peptide (C-term) - Protein Information**

Name RBBP5

Synonyms RBQ3

#### Function

In embryonic stem (ES) cells, plays a crucial role in the differentiation potential, particularly along the neural lineage, regulating gene induction and H3 'Lys-4' methylation at key developmental loci, including that mediated by retinoic acid (By similarity). Does not affect ES cell self-renewal (By similarity). Component or associated component of some histone methyltransferase complexes which regulates transcription through recruitment of those complexes to gene promoters (PubMed:<a href="http://www.uniprot.org/citations/19131338"

target="\_blank">19131338</a>). As part of the MLL1/MLL complex, involved in mono-, di- and trimethylation at 'Lys-4' of histone H3 (PubMed:<a

href="http://www.uniprot.org/citations/19556245" target="\_blank">19556245</a>). Histone H3 'Lys-4' methylation represents a specific tag for epigenetic transcriptional activation (PubMed:<a href="http://www.uniprot.org/citations/19556245" target="\_blank">19556245</a>). In association with ASH2L and WDR5, stimulates the histone methyltransferase activities of KMT2A, KMT2B, KMT2C, KMT2D, SETD1A and SETD1B (PubMed:<a



href="http://www.uniprot.org/citations/22266653" target="\_blank">22266653</a>, PubMed:<a href="http://www.uniprot.org/citations/21220120" target="\_blank">21220120</a>).

Cellular Location Nucleus.

**Tissue Location** Ubiquitously expressed.

# **RBBP5 Blocking Peptide (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

• <u>Blocking Peptides</u> RBBP5 Blocking Peptide (C-term) - Images

### RBBP5 Blocking Peptide (C-term) - Background

In embryonic stem (ES) cells, plays a crucial role in the differentiation potential, particularly along the neural lineage, regulating gene induction and H3 'Lys-4' methylation at key developmental loci, including that mediated by retinoic acid (By similarity). As part of the MLL1/MLL complex, involved in mono-, di- and trimethylation at 'Lys-4' of histone H3. Histone H3 'Lys-4' methylation represents a specific tag for epigenetic transcriptional activation.

### **RBBP5 Blocking Peptide (C-term) - References**

Saijo M., et al.Genomics 27:511-519(1995). Ota T., et al.Nat. Genet. 36:40-45(2004). Gregory S.G., et al.Nature 441:315-321(2006). Mural R.J., et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases. Hughes C.M., et al.Mol. Cell 13:587-597(2004).