

**TRRAP Blocking Peptide (C-term)**  
**Synthetic peptide**  
**Catalog # BP21568b****Specification**

---

**TRRAP Blocking Peptide (C-term) - Product Information**Primary Accession [Q9Y4A5](#)**TRRAP Blocking Peptide (C-term) - Additional Information****Gene ID** 8295**Other Names**

Transformation/transcription domain-associated protein, 350/400 kDa PCAF-associated factor, PAF350/400, STAF40, Tra1 homolog, TRRAP, PAF400

**Target/Specificity**

The synthetic peptide sequence is selected from aa 3015-3029 of HUMAN TRRAP

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**TRRAP Blocking Peptide (C-term) - Protein Information****Name** TRRAP**Synonyms** PAF400**Function**

Adapter protein, which is found in various multiprotein chromatin complexes with histone acetyltransferase activity (HAT), which gives a specific tag for epigenetic transcription activation. Component of the NuA4 histone acetyltransferase complex which is responsible for acetylation of nucleosomal histones H4 and H2A. Plays a central role in MYC transcription activation, and also participates in cell transformation by MYC. Required for p53/TP53-, E2F1- and E2F4- mediated transcription activation. Also involved in transcription activation mediated by the adenovirus E1A, a viral oncoprotein that deregulates transcription of key genes. Probably acts by linking transcription factors such as E1A, MYC or E2F1 to HAT complexes such as STAGA thereby allowing transcription activation. Probably not required in the steps following histone acetylation in processes of transcription activation. May be required for the mitotic checkpoint and normal cell cycle progression. Component of a SWR1-like complex that specifically mediates the removal of histone H2A.Z/H2AZ1 from the nucleosome. May play a role in the formation and maintenance of

the auditory system (By similarity).

#### **Cellular Location**

Nucleus

#### **TRRAP Blocking Peptide (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

#### **TRRAP Blocking Peptide (C-term) - Images**

#### **TRRAP Blocking Peptide (C-term) - Background**

Adapter protein, which is found in various multiprotein chromatin complexes with histone acetyltransferase activity (HAT), which gives a specific tag for epigenetic transcription activation. Component of the NuA4 histone acetyltransferase complex which is responsible for acetylation of nucleosomal histones H4 and H2A. Plays a central role in MYC transcription activation, and also participates in cell transformation by MYC. Required for p53/TP53-, E2F1- and E2F4-mediated transcription activation. Also involved in transcription activation mediated by the adenovirus E1A, a viral oncoprotein that deregulates transcription of key genes. Probably acts by linking transcription factors such as E1A, MYC or E2F1 to HAT complexes such as STAGA thereby allowing transcription activation. Probably not required in the steps following histone acetylation in processes of transcription activation. May be required for the mitotic checkpoint and normal cell cycle progression. Component of a SWR1- like complex that specifically mediates the removal of histone H2A.Z/H2AFZ from the nucleosome.

#### **TRRAP Blocking Peptide (C-term) - References**

McMahon S.B.,et al.Cell 94:363-374(1998).  
Vassilev A.,et al.Mol. Cell 2:869-875(1998).  
Hillier L.W.,et al.Nature 424:157-164(2003).  
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.  
Scherer S.W.,et al.Science 300:767-772(2003).