

**FLNB Blocking Peptide (N-Term)**

Synthetic peptide

Catalog # BP21863a

**Specification**

---

**FLNB Blocking Peptide (N-Term) - Product Information**

Primary Accession

[O75369](#)

Other Accession

[Q80X90](#)**FLNB Blocking Peptide (N-Term) - Additional Information**

Gene ID 2317

**Other Names**

Filamin-B, FLN-B, ABP-278, ABP-280 homolog, Actin-binding-like protein, Beta-filamin, Filamin homolog 1, Fh1, Filamin-3, Thyroid autoantigen, Truncated actin-binding protein, Truncated ABP, FLNB, FLN1L, FLN3, TABP, TAP

**Target/Specificity**

The synthetic peptide sequence is selected from aa 308-322 of HUMAN FLNB

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**FLNB Blocking Peptide (N-Term) - Protein Information**

Name FLNB

Synonyms FLN1L, FLN3, TABP, TAP

**Function**

Connects cell membrane constituents to the actin cytoskeleton. May promote orthogonal branching of actin filaments and links actin filaments to membrane glycoproteins. Anchors various transmembrane proteins to the actin cytoskeleton. Interaction with FLNA may allow neuroblast migration from the ventricular zone into the cortical plate. Various interactions and localizations of isoforms affect myotube morphology and myogenesis. Isoform 6 accelerates muscle differentiation in vitro.

**Cellular Location**

[Isoform 1]: Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, stress fiber. Cytoplasm, myofibril, sarcomere, Z line. Note=In differentiating myotubes, isoform 1, isoform 2

and isoform 3 are localized diffusely throughout the cytoplasm with regions of enrichment at the longitudinal actin stress fiber. In differentiated tubes, isoform 1 is also detected within the Z-lines [Isoform 3]: Cytoplasm, cytoskeleton, stress fiber

#### **Tissue Location**

Ubiquitous. Isoform 1 and isoform 2 are expressed in placenta, bone marrow, brain, umbilical vein endothelial cells (HUVEC), retina and skeletal muscle. Isoform 1 is predominantly expressed in prostate, uterus, liver, thyroid, stomach, lymph node, small intestine, spleen, skeletal muscle, kidney, placenta, pancreas, heart, lung, platelets, endothelial cells, megakaryocytic and erythroleukemic cell lines. Isoform 2 is predominantly expressed in spinal cord, platelet and Daudi cells. Also expressed in thyroid adenoma, neurofibrillary tangles (NFT), senile plaques in the hippocampus and cerebral cortex in Alzheimer disease (AD). Isoform 3 and isoform 6 are expressed predominantly in lung, heart, skeletal muscle, testis, spleen, thymus and leukocytes. Isoform 4 and isoform 5 are expressed in heart.

#### **FLNB Blocking Peptide (N-Term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

#### **FLNB Blocking Peptide (N-Term) - Images**

#### **FLNB Blocking Peptide (N-Term) - Background**

Connects cell membrane constituents to the actin cytoskeleton. May promote orthogonal branching of actin filaments and links actin filaments to membrane glycoproteins. Anchors various transmembrane proteins to the actin cytoskeleton. Interaction with FLNA may allow neuroblast migration from the ventricular zone into the cortical plate. Various interactions and localizations of isoforms affect myotube morphology and myogenesis. Isoform 6 accelerates muscle differentiation in vitro.

#### **FLNB Blocking Peptide (N-Term) - References**

Takafuta T.,et al.J. Biol. Chem. 273:17531-17538(1998).  
Xu W.-F.,et al.Blood 92:1268-1276(1998).  
van Der Flier A.,et al.J. Cell Biol. 156:361-376(2002).  
Chakarova C.,et al.Hum. Genet. 107:597-611(2000).  
Oshikawa M.,et al.DNA Res. 15:123-136(2008).