

CHD5 Blocking Peptide (N-Term)
Synthetic peptide
Catalog # BP22011a**Specification**

CHD5 Blocking Peptide (N-Term) - Product InformationPrimary Accession [Q8TDI0](#)**CHD5 Blocking Peptide (N-Term) - Additional Information****Gene ID** 26038**Other Names**

Chromodomain-helicase-DNA-binding protein 5, CHD-5, 3.6.4.12, ATP-dependent helicase CHD5, CHD5 {ECO:0000312|EMBL:AAL98962.1}, KIAA0444

Target/Specificity

The synthetic peptide sequence is selected from aa 84-98 of HUMAN CHD5

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

CHD5 Blocking Peptide (N-Term) - Protein Information**Name** CHD5 {ECO:0000312|EMBL:AAL98962.1}**Synonyms** KIAA0444**Function**

Chromatin-remodeling protein that binds DNA through histones and regulates gene transcription. May specifically recognize and bind trimethylated 'Lys-27' (H3K27me3) and non-methylated 'Lys-4' of histone H3. Acts as a component of the histone deacetylase NuRD complex which participates in the remodeling of chromatin. Plays a role in the development of the nervous system by activating the expression of genes promoting neuron terminal differentiation. In parallel, it may also positively regulate the trimethylation of histone H3 at 'Lys-27' thereby specifically repressing genes that promote the differentiation into non-neuronal cell lineages. Regulates the expression of genes involved in cell proliferation and differentiation. Downstream activated genes may include CDKN2A that positively regulates the p53/TP53 pathway, which in turn, prevents cell proliferation. In spermatogenesis, it probably regulates histone hyperacetylation and the replacement of histones by transition proteins in chromatin, a crucial step in the condensation of spermatid chromatin and the production of functional spermatozoa.

Cellular Location

Nucleus. Chromosome {ECO:0000250|UniProtKB:A2A8L1}

Tissue Location

Preferentially expressed in total brain, fetal brain, and cerebellum. It is also moderately expressed in the adrenal gland and detected in testis.

CHD5 Blocking Peptide (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

CHD5 Blocking Peptide (N-Term) - Images**CHD5 Blocking Peptide (N-Term) - Background**

Chromatin-remodeling protein that binds DNA through histones and regulates gene transcription. May specifically recognize and bind trimethylated 'Lys-27' (H3K27me3) and non-methylated 'Lys-4' of histone H3. Plays a role in the development of the nervous system by activating the expression of genes promoting neuron terminal differentiation. In parallel, it may also positively regulate the trimethylation of histone H3 at 'Lys-27' thereby specifically repressing genes that promote the differentiation into non-neuronal cell lineages. Tumor suppressor, it regulates the expression of genes involved in cell proliferation and differentiation. Downstream activated genes may include CDKN2A that positively regulates the p53/TP53 pathway, which in turn, prevents cell proliferation. In spermatogenesis, it probably regulates histone hyperacetylation and the replacement of histones by transition proteins in chromatin, a crucial step in the condensation of spermatid chromatin and the production of functional spermatozoa.

CHD5 Blocking Peptide (N-Term) - References

Thompson P.M., et al. *Oncogene* 22:1002-1011(2003).
Gregory S.G., et al. *Nature* 441:315-321(2006).
Bechtel S., et al. *BMC Genomics* 8:399-399(2007).
Seki N., et al. *DNA Res.* 4:345-349(1997).
Bagchi A., et al. *Cell* 128:459-475(2007).