

ARHGEF7 Blocking Peptide (C-Term) Synthetic peptide Catalog # BP22153b

Specification

ARHGEF7 Blocking Peptide (C-Term) - Product Information

Primary Accession Other Accession

<u>Q14155</u> <u>Q9ES28</u>

ARHGEF7 Blocking Peptide (C-Term) - Additional Information

Gene ID 8874

Other Names Rho guanine nucleotide exchange factor 7, Beta-Pix, COOL-1, PAK-interacting exchange factor beta, p85, ARHGEF7, COOL1, KIAA0142, P85SPR, PAK3BP, PIXB

Target/Specificity

The synthetic peptide sequence is selected from aa 674-688 of HUMAN ARHGEF7

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

ARHGEF7 Blocking Peptide (C-Term) - Protein Information

Name ARHGEF7

Synonyms COOL1, KIAA0142, P85SPR, PAK3BP, PIXB

Function

Acts as a RAC1 guanine nucleotide exchange factor (GEF) and can induce membrane ruffling. Functions in cell migration, attachment and cell spreading. Promotes targeting of RAC1 to focal adhesions (By similarity). May function as a positive regulator of apoptosis. Downstream of NMDA receptors and CaMKK-CaMK1 signaling cascade, promotes the formation of spines and synapses in hippocampal neurons.

Cellular Location

Cell junction, focal adhesion. Cell projection, ruffle. Cytoplasm, cell cortex Cell projection, lamellipodium. Note=Detected at cell adhesions. A small proportion is detected at focal adhesions



ARHGEF7 Blocking Peptide (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

Blocking Peptides

ARHGEF7 Blocking Peptide (C-Term) - Images

ARHGEF7 Blocking Peptide (C-Term) - Background

Acts as a RAC1 guanine nucleotide exchange factor (GEF) and can induce membrane ruffling. Functions in cell migration, attachment and cell spreading. Promotes targeting of RAC1 to focal adhesions (By similarity). May function as a positive regulator of apoptosis. Downstream of NMDA receptors and CaMKK-CaMK1 signaling cascade, promotes the formation of spines and synapses in hippocampal neurons.

ARHGEF7 Blocking Peptide (C-Term) - References

Nagase T., et al.DNA Res. 2:167-174(1995). Bagrodia S., et al.J. Biol. Chem. 273:23633-23636(1998). Bechtel S., et al.BMC Genomics 8:399-399(2007). Dunham A., et al.Nature 428:522-528(2004). Mural R.J., et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.