

**Ribophorin (RPN1) Antibody (C-term) Blocking peptide**  
**Synthetic peptide**  
**Catalog # BP2409b****Specification**

---

**Ribophorin (RPN1) Antibody (C-term) Blocking peptide - Product Information**Primary Accession [P04843](#)**Ribophorin (RPN1) Antibody (C-term) Blocking peptide - Additional Information**

Gene ID 6184

**Other Names**

Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit 1,  
Dolichyl-diphosphooligosaccharide--protein glycosyltransferase 67 kDa subunit, Ribophorin I,  
RPN-I, Ribophorin-1, RPN1

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP2409b](/product/products/AP2409b) was selected from the C-term region of human RPN1 . A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**Ribophorin (RPN1) Antibody (C-term) Blocking peptide - Protein Information**Name RPN1 ([HGNC:10381](#))**Function**

Subunit of the oligosaccharyl transferase (OST) complex that catalyzes the initial transfer of a defined glycan (Glc(3)Man(9)GlcNAc(2) in eukaryotes) from the lipid carrier dolichol-pyrophosphate to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent polypeptide chains, the first step in protein N-glycosylation (PubMed:[31831667](http://www.uniprot.org/citations/31831667)). N-glycosylation occurs cotranslationally and the complex associates with the Sec61 complex at the channel-forming translocon complex that mediates protein translocation across the endoplasmic reticulum (ER). All subunits are required for a maximal enzyme activity.

**Cellular Location**

Endoplasmic reticulum {ECO:0000250|UniProtKB:E2RQ08, ECO:0000250|UniProtKB:Q9GMB0}  
Endoplasmic reticulum membrane; Single-pass type I membrane protein. Melanosome.  
Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV

**Tissue Location**

Expressed in all tissues tested.

**Ribophorin (RPN1) Antibody (C-term) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**Ribophorin (RPN1) Antibody (C-term) Blocking peptide - Images****Ribophorin (RPN1) Antibody (C-term) Blocking peptide - Background**

Ribophorins 1 and 2 are abundant and highly conserved glycoproteins residing in the endoplasmic reticulum, that participate in ribosome binding. Mammalian oligosaccharyltransferase activity is associated with a protein complex composed of RPN1, RPN2, and an oligosaccharyltransferase protein. RPN1 is a component of the proteasome base. The ubiquitin-like (UBL) domain of recombinant Rad23 interacts with proteasomes through the leucine-rich repeat domain of RPN1. The RPN1 gene maps to chromosome 3 in somatic cell hybrids, and the RPN2 gene maps to chromosome 20 by in situ hybridization.

**Ribophorin (RPN1) Antibody (C-term) Blocking peptide - References**

Fu, J., et al., J. Biol. Chem. 275(6):3984-3990 (2000). Pekarsky, Y., et al., Cancer Res. 57(18):3914-3919 (1997). Crimaudo, C., et al., EMBO J. 6(1):75-82 (1987).