

**SAE1 (AOS1) Antibody (C-term) Blocking peptide**  
**Synthetic peptide**  
**Catalog # BP2511b****Specification**

---

**SAE1 (AOS1) Antibody (C-term) Blocking peptide - Product Information**Primary Accession [Q9UBE0](#)**SAE1 (AOS1) Antibody (C-term) Blocking peptide - Additional Information****Gene ID** 10055**Other Names**

SUMO-activating enzyme subunit 1, Ubiquitin-like 1-activating enzyme E1A, SUMO-activating enzyme subunit 1, N-terminally processed, SAE1, AOS1, SUA1, UBLE1A

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP2511b](/product/products/AP2511b) was selected from the C-term region of human AOS1 . A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**SAE1 (AOS1) Antibody (C-term) Blocking peptide - Protein Information****Name** SAE1**Synonyms** AOS1, SUA1, UBLE1A**Function**

The heterodimer acts as an E1 ligase for SUMO1, SUMO2, SUMO3, and probably SUMO4. It mediates ATP-dependent activation of SUMO proteins followed by formation of a thioester bond between a SUMO protein and a conserved active site cysteine residue on UBA2/SAE2.

**Cellular Location**

Nucleus.

**Tissue Location**

Expression level increases during S phase and drops in G2 phase (at protein level).

**SAE1 (AOS1) Antibody (C-term) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**SAE1 (AOS1) Antibody (C-term) Blocking peptide - Images****SAE1 (AOS1) Antibody (C-term) Blocking peptide - Background**

The dimeric enzyme AOS1 acts as a an E1 ligase for SUMO1, SUMO2, SUMO3, and probably SUMO4. It mediates ATP-dependent activation of SUMO proteins and formation of a thioester with a conserved cysteine residue on SAE2.

**SAE1 (AOS1) Antibody (C-term) Blocking peptide - References**

Desterro, J.M., et al., J. Biol. Chem. 274(15):10618-10624 (1999).Gong, L., et al., FEBS Lett. 448(1):185-189 (1999).Okuma, T., et al., Biochem. Biophys. Res. Commun. 254(3):693-698 (1999).