

PAPSS1 Antibody (N-term K9) Blocking Peptide Synthetic peptide Catalog # BP2607a

Specification

PAPSS1 Antibody (N-term K9) Blocking Peptide - Product Information

Primary Accession Other Accession

<u>043252</u> <u>NP_005434</u>

PAPSS1 Antibody (N-term K9) Blocking Peptide - Additional Information

Gene ID 9061

Other Names

Bifunctional 3'-phosphoadenosine 5'-phosphosulfate synthase 1, PAPS synthase 1, PAPSS 1, Sulfurylase kinase 1, SK 1, SK1, Sulfate adenylyltransferase, ATP-sulfurylase, Sulfate adenylate transferase, SAT, Adenylyl-sulfate kinase, 3'-phosphoadenosine-5'-phosphosulfate synthase, APS kinase, Adenosine-5'-phosphosulfate 3'-phosphotransferase, Adenylylsulfate 3'-phosphotransferase, PAPSS1, ATPSK1, PAPSS

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP2607a was selected from the N-term region of human PAPSS1 . A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

PAPSS1 Antibody (N-term K9) Blocking Peptide - Protein Information

Name PAPSS1

Synonyms ATPSK1, PAPSS

Function

Bifunctional enzyme with both ATP sulfurylase and APS kinase activity, which mediates two steps in the sulfate activation pathway. The first step is the transfer of a sulfate group to ATP to yield adenosine 5'-phosphosulfate (APS), and the second step is the transfer of a phosphate group from ATP to APS yielding 3'-phosphoadenylylsulfate (PAPS: activated sulfate donor used by sulfotransferase). In mammals, PAPS is the sole source of sulfate; APS appears to be only an



intermediate in the sulfate-activation pathway (PubMed: <a

```
href="http://www.uniprot.org/citations/9576487" target="_blank">9576487</a>, PubMed:<a
href="http://www.uniprot.org/citations/9668121" target="_blank">9668121</a>, PubMed:<a
href="http://www.uniprot.org/citations/9648242" target="_blank">9648242</a>, PubMed:<a
href="http://www.uniprot.org/citations/14747722" target="_blank">14747722</a>). Required for
normal biosynthesis of sulfated L-selectin ligands in endothelial cells (PubMed:<a
href="http://www.uniprot.org/citations/9576487" target="_blank">9576487</a>).
```

Tissue Location

Expressed in testis, pancreas, kidney, thymus, prostate, ovary, small intestine, colon, leukocytes and liver. Also expressed in high endothelial venules (HEV) cells and in cartilage

PAPSS1 Antibody (N-term K9) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

<u>Blocking Peptides</u>

PAPSS1 Antibody (N-term K9) Blocking Peptide - Images

PAPSS1 Antibody (N-term K9) Blocking Peptide - Background

Three-prime-phosphoadenosine 5-prime-phosphosulfate (PAPS) is the sulfate donor cosubstrate for all sulfotransferase (SULT) enzymes. SULTs catalyze the sulfate conjugation of many endogenous and exogenous compounds, including drugs and other xenobiotics. In humans, PAPS is synthesized from adenosine 5-prime triphosphate (ATP) and inorganic sulfate by 2 isoforms, PAPSS1 and PAPSS2.

PAPSS1 Antibody (N-term K9) Blocking Peptide - References

Venkatachalam, K.V., IUBMB Life 55(1):1-11 (2003).Xu, Z.H., et al., Biochem. Biophys. Res. Commun. 268(2):437-444 (2000).Venkatachalam, K.V., et al., J. Biol. Chem. 273(30):19311-19320 (1998).ul Haque, M.F., et al., Nat. Genet. 20(2):157-162 (1998).Girard, J.P., et al., FASEB J. 12(7):603-612 (1998).