

CCT3 Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP2890c**Specification**

CCT3 Antibody (Center) Blocking Peptide - Product InformationPrimary Accession [P49368](#)**CCT3 Antibody (Center) Blocking Peptide - Additional Information****Gene ID** 7203**Other Names**

T-complex protein 1 subunit gamma, TCP-1-gamma, CCT-gamma, hTRiC5, CCT3, CCTG, TRiC5

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP2890c](/products/AP2890c) was selected from the Center region of human CCT3. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

CCT3 Antibody (Center) Blocking Peptide - Protein Information**Name** CCT3**Synonyms** CCTG, TRiC5**Function**

Component of the chaperonin-containing T-complex (TRiC), a molecular chaperone complex that assists the folding of proteins upon ATP hydrolysis (PubMed: [25467444](http://www.uniprot.org/citations/25467444)). The TRiC complex mediates the folding of WRAP53/TCAB1, thereby regulating telomere maintenance (PubMed: [25467444](http://www.uniprot.org/citations/25467444)). As part of the TRiC complex may play a role in the assembly of BBSome, a complex involved in ciliogenesis regulating transports vesicles to the cilia (PubMed: [20080638](http://www.uniprot.org/citations/20080638)). The TRiC complex plays a role in the folding of actin and tubulin (Probable).

Cellular Location

Cytoplasm.

CCT3 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

CCT3 Antibody (Center) Blocking Peptide - Images**CCT3 Antibody (Center) Blocking Peptide - Background**

CCT3 is a molecular chaperone that is member of the chaperonin containing TCP1 complex (CCT), also known as the TCP1 ring complex (TRiC). This complex consists of two identical stacked rings, each containing eight different proteins. Unfolded polypeptides enter the central cavity of the complex and are folded in an ATP-dependent manner. The complex folds various proteins, including actin and tubulin.

CCT3 Antibody (Center) Blocking Peptide - References

Zebol,J.R., Int. J. Biochem. Cell Biol. 41 (4), 822-827 (2009)