

DLST Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP2899b**Specification**

DLST Antibody (C-term) Blocking Peptide - Product InformationPrimary Accession [P36957](#)**DLST Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 1743**Other Names**

Dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial, 2-oxoglutarate dehydrogenase complex component E2, OGDC-E2, Dihydrolipoamide succinyltransferase component of 2-oxoglutarate dehydrogenase complex, E2K, DLST, DLTS

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP2899b](/products/AP2899b) was selected from the C-term region of human DLST. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

DLST Antibody (C-term) Blocking Peptide - Protein Information**Name** DLST ([HGNC:2911](#))**Synonyms** DLTS**Function**

Dihydrolipoamide succinyltransferase (E2) component of the 2-oxoglutarate dehydrogenase complex. The 2-oxoglutarate dehydrogenase complex catalyzes the overall conversion of 2-oxoglutarate to succinyl-CoA and CO₂. The 2-oxoglutarate dehydrogenase complex is mainly active in the mitochondrion (PubMed: [29211711](http://www.uniprot.org/citations/29211711), PubMed: [30929736](http://www.uniprot.org/citations/30929736)). A fraction of the 2-oxoglutarate dehydrogenase complex also localizes in the nucleus and is required for lysine succinylation of histones: associates with KAT2A

on chromatin and provides succinyl-CoA to histone succinyltransferase KAT2A (PubMed:29211711).

Cellular Location

Mitochondrion matrix. Nucleus Note=Mainly localizes in the mitochondrion. A small fraction localizes to the nucleus, where the 2-oxoglutarate dehydrogenase complex is required for histone succinylation.

DLST Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

DLST Antibody (C-term) Blocking Peptide - Images**DLST Antibody (C-term) Blocking Peptide - Background**

DLST catalyzes the overall conversion of 2-oxoglutarate to succinyl-CoA and CO₂. It contains multiple copies of 3 enzymatic components: 2-oxoglutarate dehydrogenase (E1), dihydrolipoamide succinyltransferase (E2) and lipoamide dehydrogenase (E3).

DLST Antibody (C-term) Blocking Peptide - References

Tu,L.C., et.al., Mol. Cell Proteomics 6 (4), 575-588 (2007)