

EIF5A Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP2901c**Specification**

EIF5A Antibody (Center) Blocking Peptide - Product InformationPrimary Accession [P63241](#)**EIF5A Antibody (Center) Blocking Peptide - Additional Information****Gene ID** 1984**Other Names**

Eukaryotic translation initiation factor 5A-1, eIF-5A-1, eIF-5A1, Eukaryotic initiation factor 5A isoform 1, eIF-5A, Rev-binding factor, eIF-4D, EIF5A

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP2901c](/products/AP2901c) was selected from the Center region of human EIF5A. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

EIF5A Antibody (Center) Blocking Peptide - Protein Information**Name** EIF5A ([HGNC:3300](#))**Function**

Translation factor that promotes translation elongation and termination, particularly upon ribosome stalling at specific amino acid sequence contexts (PubMed:[33547280](http://www.uniprot.org/citations/33547280)). Binds between the exit (E) and peptidyl (P) site of the ribosome and promotes rescue of stalled ribosome: specifically required for efficient translation of polyproline-containing peptides as well as other motifs that stall the ribosome (By similarity). Acts as a ribosome quality control (RQC) cofactor by joining the RQC complex to facilitate peptidyl transfer during CAT tailing step (By similarity). Also involved in actin dynamics and cell cycle progression, mRNA decay and probably in a pathway involved in stress response and maintenance of cell wall integrity (PubMed:[16987817](http://www.uniprot.org/citations/16987817)). With syntenin SDCBP, functions as a regulator of p53/TP53 and p53/TP53-dependent apoptosis

(PubMed:15371445). Regulates also TNF-alpha-mediated apoptosis (PubMed:15452064, PubMed:17187778). Mediates effects of polyamines on neuronal process extension and survival (PubMed:17360499). Is required for autophagy by assisting the ribosome in translating the ATG3 protein at a specific amino acid sequence, the 'ASP-ASP-Gly' motif, leading to the increase of the efficiency of ATG3 translation and facilitation of LC3B lipidation and autophagosome formation (PubMed:29712776).

Cellular Location

Cytoplasm. Nucleus. Endoplasmic reticulum membrane; Peripheral membrane protein; Cytoplasmic side. Note=Hypusine modification promotes the nuclear export and cytoplasmic localization and there was a dynamic shift in the localization from predominantly cytoplasmic to primarily nuclear under apoptotic inducing conditions (PubMed:19379712, PubMed:27306458). Nuclear export of hypusinated protein is mediated by XPO4 (PubMed:10944119, PubMed:27306458).

Tissue Location

Expressed in umbilical vein endothelial cells and several cancer cell lines (at protein level)

EIF5A Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

EIF5A Antibody (Center) Blocking Peptide - Images

EIF5A Antibody (Center) Blocking Peptide - Background

EIF5A precursor is the only cellular protein known to contain a specific lysine residue which is transformed into the unique amino acid hypusine [N-(4-amino-2-hydroxybutyl)-lysine] by a series of post translational reactions. eIF5A promotes the formation of the first peptide bond during the initial stage of protein synthesis.

EIF5A Antibody (Center) Blocking Peptide - References

Lee,S.B., et.al., Biochem. Biophys. Res. Commun. 383 (4), 497-502 (2009)Gosslau,A., et.al., J. Cell. Physiol. 219 (2), 485-493 (2009)