

# PGAM1 Antibody (Center) Blocking Peptide

Synthetic peptide Catalog # BP2923c

#### Specification

## PGAM1 Antibody (Center) Blocking Peptide - Product Information

Primary Accession

#### <u>P18669</u>

## PGAM1 Antibody (Center) Blocking Peptide - Additional Information

Gene ID 5223

## **Other Names** Phosphoglycerate mutase 1, BPG-dependent PGAM 1, Phosphoglycerate mutase isozyme B, PGAM-B, PGAM1, PGAMA

## Target/Specificity

The synthetic peptide sequence used to generate the antibody <a href=/products/AP2923c>AP2923c</a> was selected from the Center region of human PGAM1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions** This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## PGAM1 Antibody (Center) Blocking Peptide - Protein Information

Name PGAM1 (HGNC:8888)

#### Synonyms PGAMA

Function

Catalyzes the interconversion of 2-phosphoglycerate and 3- phosphoglyceratea crucial step in glycolysis, by using 2,3- bisphosphoglycerate (PubMed:<a href="http://www.uniprot.org/citations/23653202" target="\_blank">23653202</a>). Also catalyzes the interconversion of (2R)-2,3-bisphosphoglycerate and (2R)-3-phospho- glyceroyl phosphate (PubMed:<a href="http://www.uniprot.org/citations/23653202" target="\_blank">23653202</a>). Also catalyzes the interconversion of (2R)-2,3-bisphosphoglycerate and (2R)-3-phospho- glyceroyl phosphate (PubMed:<a href="http://www.uniprot.org/citations/23653202" target="\_blank">23653202" target="\_blank">23653202</a>).

**Tissue Location** 

Expressed in the liver and brain. Not found in the muscle.



## PGAM1 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

#### <u>Blocking Peptides</u>

## PGAM1 Antibody (Center) Blocking Peptide - Images

#### PGAM1 Antibody (Center) Blocking Peptide - Background

PGAM1 can be interconversion of 3- and 2-phosphoglycerate with 2,3-bisphosphoglycerate as the primer of the reaction. It can also catalyze the reaction of EC 5.4.2.4 (synthase) and EC 3.1.3.13 (phosphatase), but with a reduced activity.

#### **PGAM1 Antibody (Center) Blocking Peptide - References**

Martins-de-Souza, D., et.al., BMC Psychiatry 9, 17 (2009)Oh, S.J., et.al., Muscle Nerve 34 (5), 572-576 (2006)